

Eulerian, Lagrangian and Broad continuous solutions to a balance law with non-convex flux. I

GIOVANNI ALBERTI, STEFANO BIANCHINI, LAURA CARAVENNA

ABSTRACT. We discuss different notions of *continuous* solutions to the balance law

$$\partial_t u + \partial_x(f(u)) = g$$

with g bounded and $f \in C^2$, extending previous works relative to the flux $f(u) = u^2$. We establish the equivalence among distributional solutions and a suitable notion of Lagrangian solutions for general smooth fluxes. We eventually find that continuous solutions are Kruřkov iso-entropy solutions, which yields uniqueness for the Cauchy problem. We also reduce the ODE on *any* characteristics under the sharp assumption that the set of inflection points of the flux f is negligible. The correspondence of the source terms in the two settings is matter of the companion work [2], where we include counterexamples when the negligibility on inflection points fails.

KEYWORDS: balance laws, Lagrangian description, Eulerian formulation.

MSC (2010): 35L60, 37C10, 58J45.

CONTENTS

1. Introduction	2
1.1. Definitions and setting	3
1.2. Overview of the results	7
2. Lagrangian solutions are distributional solutions	8
2.1. The case of <i>BV</i> -regularity	9
2.2. The case of continuous solutions: <i>BV</i> approximations	14
3. Distributional solutions are broad solutions, if inflections are negligible	20
3.1. Lipschitz regularity along characteristics	20
3.2. Construction of a universal source	22
4. Distributional solutions are Lagrangian solutions	30
Appendix A. Three sufficient conditions for the Lagrangian formulation	35
A.1. A dense set of characteristics	35
A.2. Lipschitz continuity along characteristics	36
A.3. Stability of the Lagrangian formulation for uniform convergence of u	37
Nomenclature	38
Acknowledgements	39
References	39

1. INTRODUCTION

Single balance laws in one space dimension mostly present smooth fluxes, although the case of piecewise smooth fluxes is of interest both for the mathematics and for applications. Source terms instead are naturally rough, and singularities of different nature have a physical and geometrical meaning. As well, they might indeed make a difference among the Eulerian and Lagrangian description of the phenomenon which is being modeled, for the mathematics.

We are concerned in this paper with different notions of *continuous* solutions of the PDE

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g(t, x), \quad (1.1)$$

where $f \in C^2(\mathbb{R})$ and the source term g is bounded. An essential feature of conservation laws is that solutions to the Cauchy problem do develop shocks in finite time. Nevertheless, the source might act as a control device for preventing this shock formation: exploiting the geometric interplay and correspondence with intrinsic Lipschitz surfaces in the Heisenberg group, [11, 5] show, for the quadratic flux, that the Cauchy problem admits continuous solutions for any Hölder continuous initial datum, if one chooses accordingly a bounded source term.

This framework of continuous solutions, with more regularity assumptions on the source term, was already considered in [8] as the natural class of solutions to certain interesting dispersive partial differential equations that can be recast as balance laws. We believe that our study is relevant also in order to point out that, even in the analysis of a single equation in one space dimension, the mathematical difficulties do not only arise by the presence of shocks: also the study of continuous solutions has important delicate points which are not technicalities. This fails the expectation that the study of continuous solutions should be easy, and equation (1.1) is a toy-model for more complex situations.

One can adopt the Eulerian viewpoint or the Lagrangian/Broad viewpoint: roughly, the first interprets the equation in a distributional sense while the second consists in an infinite dimensional system of ODEs along characteristics. We compare here the equivalence among the formulations when u is assumed to be continuous, but no more. We remark that even with the quadratic flux

$$f(z) = z^2/2$$

in general u is not more than Hölder continuous, see [12], so that a finer analysis is needed. Continuous solutions are regularized to locally Lipschitz on the open set $\{f'(u)f''(u) \neq 0\}$, exploiting the results of this paper, for time-dependent solutions when the source term g is autonomous [1], but not in general. Examples of stationary solutions which are neither absolutely continuous nor of bounded variation are trivially given by continuous functions $u(x)$ for which $f(u(x))$ has bounded derivative.

Correspondences among different formulations are already done at different levels in [8, 19, 4, 5, 7] for the special but relevant case of the quadratic flux. We extend the analysis with new tools. The issue is delicate because g in this setting lacks even of continuity in the x variable, and characteristic curves need not be unique because u lacks of smoothness. As a consequence, the source terms for the two descriptions lie in different spaces:

- in the Eulerian point of view g is identified only as a distribution in the (t, x) -space;
- in the Lagrangian/Broad viewpoint it is the restriction of g on any characteristic curve which must identify uniquely a distribution in the t -space—or for a weaker notion only on a chosen family of characteristics that we call Lagrangian parameterization.

The aim of this paper is to consider and to discuss when Eulerian, Broad and Lagrangian solution of (1.1) that we just mentioned are equivalent notions, without addressing what is the correspondence among the suitable source terms—if any. The correspondence of the source terms, source terms which belong to different functional spaces, is the subject of the companion paper [2], including counterexamples which show that the formulations are not always equivalent.

We conclude mentioning that Broad solutions were introduced in [17] as generalizations of classical solutions alternative to the distributional (Eulerian) ones, and presented e.g. also in [6]. They were successfully studied and applied in different situations where characteristic curves are unique; the analysis in situations when characteristics do merge and split however was only associated to the presence of shocks, and a different analysis related to multi-valued solutions was performed. They were then considered for the quadratic flux by F. Bigolin and F. Serra Casano for their interest related to intrinsic regular and intrinsic Lipschitz surfaces in the Heisenberg group. Our notions of Lagrangian and Broad solutions collapse and substantially coincide with the ones in the literature when the settings overlap. They are otherwise a nontrivial extension of those concepts, and most of the issues in the analysis arise because of our different setting.

1.1. Definitions and setting

Since we are in a non-standard setting, we explain extensively the different notions of solutions and we specify the notation we adopt. Even if this is an heavy block, detailed definitions improve the later analysis. They will be also collected in the Nomenclature at the end for an easy consultation.

Notation 1. We can assume below that $u \in C_c(\mathbb{R}^+ \times \mathbb{R})$, because our considerations are local in space-time. We adopt the short notation $\lambda(t, x) = f'(u(t, x))$ for the characteristic speed.

Notation 2. Given a function of two variables $\varphi(t, x)$, one denotes the restrictions to coordinate sections as

$$\varphi_x^{e1} : t \mapsto \varphi(t, x), \quad \varphi_t^{e2} : x \mapsto \varphi(t, x).$$

Notation 3. Given a function of locally bounded variation $\varphi(t, x)$, one denotes by

$$D_t \varphi(dt, dx), \quad D_x \varphi(dt, dx)$$

the measures of its partial derivatives. When it is not known if they are measures, we rather denote the distributional partial derivatives by

$$\partial_t \varphi(t, x), \quad \partial_x \varphi(t, x).$$

Classical partial derivatives are often denoted by

$$\frac{\partial \varphi(t, x)}{\partial t}, \quad \frac{\partial \varphi(t, x)}{\partial x}.$$

Definition 4 (Characteristic Curves). *Characteristic curves* of u are absolutely continuous functions $\gamma : \mathbb{R}^+ \rightarrow \mathbb{R}$, or equivalently the corresponding curves $i_\gamma(s) := (s, \gamma(s))$, defined on a connected open subset of \mathbb{R} and satisfying the ordinary differential equation

$$\dot{\gamma}(s) = \lambda(s, \gamma(s)) = \lambda(i_\gamma(s)).$$

The continuity of u implies that γ is continuously differentiable.

Notice that i_γ is an integral curve of the vector field $(1, \lambda)$.




FIGURE 1. Curves satisfying $\dot{\gamma}(s) = 1$ almost everywhere might fail to be Lipschitz continuous: characteristic curves are indeed required to be absolutely continuous. Absolute continuity automatically improves to C^1 -regularity, due to the continuity of the field. They are then stable under uniform convergence.

Definition 5 (Lagrangian Parameterization). We call *Lagrangian parameterization* associated with u a surjective continuous function $\chi : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$, or equivalently $\chi : \mathbb{R} \rightarrow C(\mathbb{R}^+)$, such that¹

- for each $y \in \mathbb{R}$, the curve $\chi(y)$ defined by $t \mapsto \chi(t, y) = \chi_y^{e_1}(t)$ is a characteristic curve:

$$\dot{\chi}_y^{e_1}(t) = \partial_t \chi(t, y) = f'(u(t, \chi(t, y))) = \lambda(t, \chi(t, y)) = \lambda(i_{\chi(y)}(t));$$

- for each $t \in \mathbb{R}^+$, $y \mapsto \chi(t, y) = \chi_t^{e_2}(y)$ is nondecreasing.

¹ There is no reason for asking the following condition only for \mathcal{L}^1 -a.e. y : if it holds for \mathcal{L}^1 -a.e. y then it holds naturally for every y . As well, it would be odd requiring the second condition only \mathcal{L}^1 -a.e. t .

Definition 6. We call a Lagrangian parameterization χ *absolutely continuous* if $(i_\chi^{-1})_\# \mathcal{L}^2 \ll \mathcal{L}^2$. Equivalently, \mathcal{L}^2 -positive measure sets can not be negligible along the characteristics of the parameterization χ : χ maps negligible sets into negligible sets.

Remark 7. Even if χ generally is not injective, $(i_\chi^{-1})_\# \mathcal{L}^2$ is as well a well defined Borel measure meaning that for all compact subsets $K \subset \mathbb{R}^+ \times \mathbb{R}$ one defines $(i_\chi^{-1})_\# \mathcal{L}^2 := \mathcal{L}^2(i_\chi(K))$. Of course $((i_\chi^{-1})_\# \mathcal{L}^2)(\emptyset) = 0$ but one has also that if A and B are disjoint compact subsets of the plane then for all t the intersection of $\chi(A \cap \{t\})$ and $\chi(B \cap \{t\})$ is at most countable, due to the monotonicity of $\chi_t^{e_2}$:

$$\mathcal{L}^1(\chi(A \cap \{t\}) \cap \chi(B \cap \{t\})) = 0 \text{ for every } t \Rightarrow \mathcal{L}^2(i_\chi(A) \cap i_\chi(B)) = 0.$$

In particular

$$\mathcal{L}^2(i_\chi(A)) + \mathcal{L}^2(i_\chi(B)) = \mathcal{L}^2(i_\chi(A) \cup i_\chi(B)) = \mathcal{L}^2(i_\chi(A \cup B)).$$

This implies that $(i_\chi^{-1})_\# \mathcal{L}^2$ is countably-additive, and thus a measure. This justifies our notation.

Notation 8. We fix the following nomenclature, that we extend at the end of the paper:

- X, k usually X is a subset $\mathbb{R}^+ \times \mathbb{R}$ and $k \in \mathbb{N} \cup \{\infty\}$;
- Ω open subset of $\mathbb{R}^+ \times \mathbb{R}$, usually it is also supposed to be bounded;
- $C(X)$ space of continuous functions on X ;
- $C_b(X)$ space of bounded continuous functions on X ;
- $C^{1/\alpha}(X)$ space of $1/\alpha$ -Hölder continuous functions on X , where $0 < 1/\alpha \leq 1$;
- $C_{(c)}^k(\Omega)$ space of k -times continuously differentiable (compactly supported) functions on Ω ;
- $C^{k,1/\alpha}(\Omega)$ space of k -times continuously differentiable (compactly supported) functions on Ω whose k -th derivative which is $1/\alpha$ -Hölder continuous in Ω ;
- $\mathfrak{L}^\infty(X)$ space of bounded Borel functions \mathfrak{g} on X ;
- $\mathfrak{L}_B^\infty(X)$ space of equivalence classes \mathfrak{g}_B made of those bounded Borel functions which coincide \mathcal{L}^1 -a.e. when restricted to *any* characteristic curve of u ;
- $\mathfrak{L}_L^\infty(X)$ space of equivalence classes \mathfrak{g}_L made of those bounded Borel functions which coincide \mathcal{L}^1 -a.e. when restricted to $\{i_{\chi(y)}(t)\}_{t>0}$, for every $y \in \mathbb{R}$ and for a fixed Lagrangian parameterization χ ;
- $L^\infty(X)$ space of equivalence classes \mathfrak{g}_E of Borel bounded functions which coincide \mathcal{L}^2 -a.e.
- $\mathcal{D}(\Omega)$ space of distributions on Ω ;
- $\mathcal{M}(X)$ space of Radon measures on X .

Notation 9. Notice there are the following natural correspondences

$$\begin{array}{ccc} \mathfrak{L}^\infty(X) & \xrightarrow{[\cdot]_\chi} & \mathfrak{L}_B^\infty(X) & \xrightarrow{[\cdot]_\chi} & \mathfrak{L}_L^\infty(X) \\ \mathfrak{g} & \mapsto & \mathfrak{g}_B = [\mathfrak{g}]_\lambda & \mapsto & \mathfrak{g}_L = [\mathfrak{g}]_\chi \end{array}$$

and moreover

$$\begin{array}{ccc} \mathfrak{L}^\infty(X) & \xrightarrow{[\cdot]} & L^\infty(X) \\ \mathfrak{g} & \mapsto & g_E = [\mathfrak{g}] \end{array}$$

The same brackets denote also correspondences from any of the bigger spaces: brackets identify the target spaces. The correspondences among $\mathfrak{L}_L^\infty(X)$, $\mathfrak{L}_B^\infty(X)$ and $L^\infty(X)$ do not exist in general. Trivially, sets which are \mathcal{L}^2 -negligible generally are not \mathcal{L}^1 -negligible along any characteristic curve γ of (1.1). Moreover, there exists a subset of the plane which has positive Lebesgue measure but which intersects each characteristic curve of a Lagrangian parameterization in a single point. See [2, §4.1-2]. A correspondence exists with *absolute continuity*.

Lemma 10. *If a Lagrangian parameterization χ is absolutely continuous, for every Borel functions $\mathfrak{g}_1, \mathfrak{g}_2 \in \mathfrak{L}^\infty(X)$ such that $[\mathfrak{g}_1]_\chi = [\mathfrak{g}_2]_\chi$ one has that $[\mathfrak{g}_1] = [\mathfrak{g}_2] \in L^\infty(X)$.*

Proof. It is an algebraic application of the definitions of the spaces in Notation 8. \square

Definition 11 (Distributional solution). Let $g_E \in L^\infty(\Omega)$. If $u \in C(\Omega)$ satisfies

$$\iint_{\Omega} \varphi_t u + \varphi_x f(u) = \iint_{\Omega} \varphi g_E \quad \forall \varphi \in C_c^\infty(\Omega)$$

we say that u is a *continuous distributional (or Eulerian) solution* of (1.1).

Definition 12 (Lagrangian solution). A function $u \in C(\Omega)$ is called a *continuous Lagrangian solution* of (1.1) with Lagrangian parameterization χ associated with u , and Lagrangian source term $g_L \in \mathfrak{L}_L^\infty(\Omega)$, if

$$\frac{d}{dt}u(t, \chi(t, y)) = g_L(t, \chi(t, y)) \quad \text{in } \mathcal{D}(i_{\chi(y)}^{-1}(\Omega)) \text{ for all } y \in \mathbb{R}.$$

Definition 13 (Broad solution). Let $u \in C(\Omega)$ and $g_B \in \mathfrak{L}_B^\infty(\Omega)$. The function u is called a *continuous broad solution* of (1.1) if it satisfies

$$\frac{d}{dt}u(t, \gamma(t)) = g_B(t, \gamma(t)) \quad \text{in } \mathcal{D}(i_\gamma^{-1}(\Omega)) \text{ for all characteristic curves } \gamma \text{ of } u.$$

Definition 14. A continuous function u is both a distributional/Lagrangian/broad solution of (1.1) when the source terms are *compatible*: if there exists a Borel function \mathfrak{g} such that

$$g_E = [\mathfrak{g}], \quad g_L = [\mathfrak{g}]_\chi, \quad g_B = [\mathfrak{g}]_\lambda.$$

Definition 15. We define $z^* \in \mathbb{R}$ an *inflection point* of a function $f \in C^2(\mathbb{R})$ if $f''(z^*) = 0$, but z^* is neither a local maximum nor a local minimum for the function $z \mapsto f(z) - f'(z^*)(z - z^*)$. We denote by $\text{Infl}(f)$ the set of inflection points of f , and by $\text{clos}(\text{Infl}(f))$ its closure.

In principle, u could be a distributional solution of (1.1) with source g_E and a Lagrangian solution with source g_L with g_E and g_L which do not correspond to a same function $\mathfrak{g} \in \mathfrak{L}^\infty(\mathbb{R}^+ \times \mathbb{R})$: in this case, we would not say that u is both a

distributional and Lagrangian solution to the *same* equation, because source terms are different. We discuss the issue in [2], where we prove that if the inflection points of f are negligible then whenever a same function is a Lagrangian solution and it is a distributional solution then the source terms must be compatible.

1.2. Overview of the results

Now that the definitions are clear, we describe our results: in §1.2.1 we collect the observations on Lagrangian parameterizations and Lagrangian/Broad solutions, and in §1.2.2 we summarize the relations among the different notions of solutions of (1.1).

Notice first that both the definitions above and the statements below are local in space-time, as well as the compatibility of the sources that will be discussed in [2]. This motivates the assumption that u is compactly supported, that we fixed in Notation 1.

1.2.1. Auxiliary observations

We begin collecting elementary observations on the basic concept of Lagrangian parameterization and Lagrangian/Broad solution, mostly for consistency.

Lemma 16. *There exists a Lagrangian parameterization χ associated with any $u \in C_b(\Omega)$. In particular, one has the implication*

$$\text{continuous broad solution} \Rightarrow \text{continuous Lagrangian solution with } g_L = [g_B]_\chi.$$

The converse implication holds under a condition on the inflection points of f , not in general.

Proof. An explicit construction of a Lagrangian parameterization χ is part of §A.1. It relies on Peano's existence theorem for ODEs with continuous coefficients. If $g_B \in \mathfrak{L}_B^\infty(\Omega)$ is the Broad source, then $[g_B]_\chi$ is immediately the Lagrangian source. The converse implication does not always hold, see [2, §4.3]. \square

Lemma 17. *Let $u \in C(\text{clos } \Omega)$ and $G > 0$. Assume that through every point of a dense subset of Ω there exists a characteristic curve along which u is G -Lipschitz continuous. Then there exists a Lagrangian parameterization χ along whose characteristics u is G -Lipschitz continuous.*

The proof of this lemma is given in §A.1.

Lemma 18. *Let $u \in C(\text{clos}(\Omega))$ and $G > 0$. A sufficient condition for u being a Lagrangian solution of (1.1) is the existence of a Lagrangian parameterization χ such that*

$$\text{the distributions } \frac{d}{dt}u(t, \chi(t, y)) \text{ are bounded in } \mathcal{D}(i_{\chi(y)}^{-1}(\Omega)) \text{ uniformly in } y.$$

The proof of this lemma is given in §A.2.

1.2.2. Main results

In the present paper we do not discuss existence of continuous solutions of (1.1), but we assume that we are given a continuous function u : due to the lack of regularity, the focus of this paper is in which sense it can be a solution of (1.1).

We first state one of the important conditions: we denote by (H) the assumption:

the set $\text{clos}(\text{Infl}(f))$ (see Definition 15) is \mathcal{L}^1 -negligible. (H)

Our results are *roughly* summarized by the following implications:

$$\text{Broad} \begin{array}{c} \Rightarrow (\text{Th. 16}) \\ \Leftarrow \text{if (H) holds } (\S 3) \end{array} \text{Lagrangian} \begin{array}{c} \Rightarrow (\S 2) \\ \Leftarrow (\S 4) \end{array} \text{distributional.}$$

The distinction among Lagrangian and distributional continuous solutions is motivated by the fact that the two formulations are different, and it is not that trivial proving their equivalence. Moreover, Lagrangian and distributional source terms do not correspond automatically, as we discuss in [2]. In particular, if we do not assume the negligibility of inflection points we are not yet able to say that the Lagrangian and distributional source terms must be compatible. If, for example, the flux function is analytic, then our work gives a full analysis.

We collect in the table below interesting properties of the solution. These properties depend on general assumptions on the smooth flux function f :

- whether f satisfies a convexity assumption named in [2] α -convexity, $\alpha > 1$, which for $\alpha = 2$ is the classical uniform convexity;
- whether the closure of inflection points of f is negligible, as in (H) above.

	α -convexity	negligible inflections	general case
absolutely continuous Lagrangian parametrization	✗ [2, §4.1]	✗	✗
u Hölder continuous	✓ [2, §2.1]	✗ [2, §4.2]	✗
u \mathcal{L}^2 -a.e. differentiable along characteristic curves	✓ [2, §2.2]	✗ [2, §4.2]	✗
u Lipschitz continuous along characteristic curves	✓	✓ Theorem 30	✗ [2, §4.3]
entropy equality	✓	✓	✓ Lemma 42
compatibility of sources	✓✓ [2, §2.2]	✓ [2, §3]	

We show in Corollary 21 that if the continuous solution u has bounded total variation then one can as well select a Lagrangian parameterization which is absolutely continuous, for $f \in C^2$.

2. LAGRANGIAN SOLUTIONS ARE DISTRIBUTIONAL SOLUTIONS

Consider a continuous Lagrangian solution $u(t, x)$ of (1.1) in the sense of Definition 12. Let χ be a Lagrangian parameterization, $g_L \in \mathcal{L}_L^\infty(\Omega)$ be its source term and set $G = \|g_L\|_\infty$. We want to show that there exists $g_E \in L^\infty(\Omega)$ with $|g_E(t, x)| \leq G$ such that $u(t, x)$ is a distributional solution of

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g_E(t, x). \quad (1.1)$$

We do not discuss at this stage the compatibility of the source terms g_L and g_E .

Notation 19. We already observed in the introduction that we are considering local statements. We directly assume therefore

- $\Omega = \mathbb{R}^+ \times \mathbb{R}$;
- u compactly supported.

We set $\Lambda = \max \lambda = \max f'(u)$. Recall that we set $G = \|g_L\|_\infty$.

2.1. The case of BV-regularity

In the present section we assume that u is not only continuous but also that it has bounded variation. Under this simplifying assumption, we prove in Lemma 22 below that u is a distributional solution to (1.1), with the natural candidate for g_E . The proof is based on explicit computations. Computations of this section exploit Vol'pert chain rule and the possibility to produce a change of variables which is absolutely continuous, as we state in Corollary 21 below. It follows by the following more general lemma.

Lemma 20. Consider a function $w : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$ such that

- the restriction w_y^{e1} belongs to $C^{1,1}(\mathbb{R}^+)$ for all $y \in \mathbb{R}$;
- the second mixed derivative $\partial_{ty}w$ is a Radon measure.

Then, up to re-parametrizing the y -variable, there exists $0 \leq H \in L_{\text{loc}}^\infty(\mathbb{R}^+ \times \mathbb{R})$ such that

$$D_y w = H \mathcal{L}^2, \quad D_t H = D_y \left(\frac{\partial w}{\partial t} \right) \in \mathcal{M}(\mathbb{R}^+ \times \mathbb{R}).$$

We rather prefer to prove the following corollary, which is more related to the notation we adopt: the proof of Lemma 20 is entirely analogous. The irrelevant disadvantage is that the commutation of the t - and y - distributional derivatives is less evident than in the above lemma.

Corollary 21. Let u be a continuous Lagrangian solution of (1.1) such that $\partial_x u(t, x)$ is a Radon measure. Then one can choose a Lagrangian parameterization χ which is absolutely continuous (see Definition 6): this additional regularity allows the injection

$$\begin{array}{ccc} \mathcal{L}_L^\infty(\Omega) & \xrightarrow{[\cdot]} & L^\infty(\Omega), \\ g_L = [g]_\chi & \mapsto & g_E = [g] = [g_L]. \end{array}$$

Moreover, for every test function $\Phi(t, y) \in C_c^1(\mathbb{R}^+ \times \mathbb{R})$ and for \mathcal{L}^1 -a.e. t one has

$$\frac{d}{dt} \int \Phi(t, y) H(t, y) dy = \int \Phi(t, y) Df'(U_t^{\text{e1}}(dy)) + \int \frac{\partial \Phi(t, y)}{\partial t} H(t, y) dy. \quad (2.1)$$

Lemma 22. Under the assumptions of Corollary 21, u has locally bounded variation. Moreover, denoting by $g_L = [g]_\chi$ a Lagrangian source, then one has

$$D_t u(dt, dx) + D_x f(u(dt, dx)) = g(t, x) dt dx.$$

Note that Lemma 21 does not follow from the theory of ODEs with rough coefficients because the notion of Lagrangian parameterization is more specific than a solution of a system of ODEs. In this paper, where the focus is on the PDE (1.1), we rather prefer to prove the lemma in the form of Corollary 21. We stress once more that computations below, switching notations, prove indeed Lemma 20, proof which could be slightly shortened in a more abstract setting.

Remark 23. Let $U(t, y) := u(t, \chi(t, y))$ for some Lagrangian parameterization χ . We notice that one can equivalently assume that either $\partial_x u$ or $\partial_y U(t, y)$ is a Radon measure. This is a direct consequence of the slicing theory of BV functions, because $\chi_t^{\text{e}2}(y)$ is monotone and the total variation of $u_t^{\text{e}2}(x)$ is equal to the total variation of $U_t^{\text{e}2}(y)$.

Proof of Lemma 21. Let χ be a Lagrangian parameterization corresponding to u . By assumption and by Remark 23, for \mathcal{L}^1 -a.e. t also the function

$$y \mapsto u(i_{\chi(y)}(t)) =: U_t^{\text{e}1}(y)$$

has locally bounded variation. Moreover, for every y by definition of Lagrangian solution

$$t \mapsto u(i_{\chi(y)}(t)) =: U_y^{\text{e}2}(t)$$

is Lipschitz continuous. We deduce by the slicing theory of BV functions [3, Th. 3.103] that also the function $(t, y) \mapsto U(t, y)$ has locally bounded variation. We show now that the Lagrangian parameterization χ can be here assumed to be absolutely continuous.

Step 1. Renormalization of y for absolute continuity of χ . Consider the two coordinate disintegrations of the measure on the plane given by $D_y U(t, y)$: by the classical disintegration theorem [3, Th. 2.28] there exists a nonnegative Borel measure $m \in \mathcal{M}^+(\mathbb{R})$ and a measurable measure-valued map $y \mapsto \nu_y \in \mathcal{M}^+(\mathbb{R})$ such that

$$D_y U(dt, dy) = \int DU_t^{\text{e}1}(dy) dt = \int \nu_y(dt) m(dy). \quad (2.2)$$

The first equality is just the slicing theory for BV functions [3, Th. 3.107].

Claim 24. Consider the Lagrangian parameterization $\bar{\chi}(t, y) := \chi(t, h^{-1}(y))$ with h defined by

$$h(y) := y + m((-\infty, y]) + D\chi_0^{\text{e}2}((-\infty, y]).$$

Then one has that $h_{\#}m \ll \mathcal{L}^1$ and $h_{\#}D\chi_0^{\text{e}2} \ll \mathcal{L}^1$ with densities bounded by 1.

Proof of Claim 24. Fix any $a \leq b$. We first observe that

$$\begin{aligned} h(b) - h(a) &= b - a + m((a, b]) + D\chi_0^{\text{e}2}((a, b]) \\ &\geq \max\{b - a, m((a, b]), D\chi_0^{\text{e}2}((a, b])\} \geq 0. \end{aligned}$$

This shows that $Dh \geq \mathcal{L}^1$, $Dh \geq m$ and $Dh \geq D\chi_0^{\text{e}2}$. Since U and χ are continuous functions, then $D_y U$ and $D\chi_0^{\text{e}2}$ are continuous measures and therefore h is a continuous function. Fix any $a < b$ and suppose $h(a') = a$, $h(b') = b$. Then one verifies

that

$$\begin{aligned} \frac{h_{\#}m([a, b])}{b - a} &\leq \frac{m([a', b'])}{h(b') - h(a')} \\ &\leq \frac{m([a', b'])}{b' - a' + m((a', b']) + D\chi_0^{\text{e}2}((a', b'))} < 1. \end{aligned}$$

This concludes the proof for $h_{\#}m$, and $h_{\#}D\chi_0^{\text{e}2}$ is entirely similar. \square

Claim 24 assures that one can re-parametrize the y -variable so that both $m(dy)$ and $D\chi_0^{\text{e}2}(y)$ are absolutely continuous with bounded densities. Let $\vartheta(y)$ and $\beta(y)$ be their Radon–Nikodym derivatives w.r.t. \mathcal{L}^1 after, eventually, the re-parametrization of y :

$$m(dy) =: \vartheta(y) dy, \quad D\chi_0^{\text{e}2}(y) =: \beta(y) dy, \quad \vartheta, \beta \in L^\infty. \quad (2.3)$$

Step 2. Formula for $D_y \chi$. In this step we prove Claim 25 below, which implies that both the distributional partial derivatives of χ are absolutely continuous measures. The claim below yields then that χ is an absolutely continuous Lagrangian parameterization. As $\chi(t, y)$ maps negligible sets into negligible sets, then one has the inclusion stated in Lemma 10:

$$\mathcal{L}_L^\infty(\mathbb{R}^+ \times \mathbb{R}) \hookrightarrow L^\infty(\mathbb{R}^+ \times \mathbb{R}).$$

Claim 25. The measure $D_y \chi(dt, dy)$ is given by the following formula:

$$D_y \chi(dt, dy) = H(t, y) dt dy, \quad L_{\text{loc}}^\infty(i_\chi^{-1}(\Omega)) \ni H(t, y) \geq 0 \quad (2.4a)$$

$$H(t, y) = \vartheta(y) \int_0^t f''(U(s, y)) \nu_y(ds) + \beta(y). \quad (2.4b)$$

Proof of Claim 25. Being

$$\dot{\chi}_y^{\text{e}1}(t) = f'(U(t, y)) \quad (2.5)$$

by Vol'pert chain rule [3, Th. 3.96] and (2.2) one has the following disintegration

$$\begin{aligned} D_y f'(U(dt, dy)) &= f''(U(t, y)) D_y U(dt, dy) \\ &= \int \{f''(U(t, y)) \nu_y(dt)\} m(dy). \end{aligned} \quad (2.6)$$

One can compute $D_y \chi$ in the following way: write χ as a primitive and differentiate under the integral. For every test function $\Phi(t, y)$

$$\begin{aligned}
& - \iint \Phi(t, y) D_y \chi(dt, dy) = \\
& = \iint \frac{\partial \Phi(t, y)}{\partial y} \chi(t, y) dy dt \\
& = \iint \frac{\partial \Phi(t, y)}{\partial y} \left\{ \int_0^t \dot{\chi}_y^{e_1}(s) ds + \chi(0, y) \right\} dy dt \\
& = \iint \int_0^t \frac{\partial \Phi(t, y)}{\partial y} [\dot{\chi}_y^{e_1}(s) + \dot{\chi}_0^{e_2}(y)/t] ds dy dt \\
& \stackrel{(2.5)}{=} \int \left\{ \iint_0^t \frac{\partial \Phi(t, y)}{\partial y} [f'(U(s, y)) + \dot{\chi}_0^{e_2}(y)/t] ds dy \right\} dt \\
& = - \int \left\{ \iint_0^t \Phi(t, y) [D_y f'(U(ds, dy)) + D \dot{\chi}_0^{e_2}(y)/t] \right\} dt
\end{aligned}$$

The last step was allowed because $U \in BV$. Owing to (2.6), (2.3) we can now proceed with

$$\begin{aligned}
& = - \int \left\{ \iint_0^t \Phi(t, y) f''(U(s, y)) \nu_y(ds) \vartheta(y) dy \right\} dt \\
& \quad - \int \left\{ \iint_0^t [\beta(y)/t] ds dy \right\} dt \\
& = - \iint \left\{ \Phi(t, y) \left[\beta(y) + \vartheta(y) \int_0^t f''(U(s, y)) \nu_y(ds) \right] \right\} dy dt.
\end{aligned}$$

Note that H is the function within the inner square brackets, thus we proved the claim. \square

Step 3. Time derivative of H . Definition (2.4b) of H does not directly allow to differentiate H in the t variable, because the measure ν_y may not be absolutely continuous. Nevertheless, this is possible from (2.4a), obtaining that $D_t H$ is a Radon measure.

Claim 26. For every test function $\Phi(t, y) \in C_c^1(\mathbb{R}^+ \times \mathbb{R})$ and for \mathcal{L}^1 -a.e. t one has (2.1)

$$\frac{d}{dt} \int \Phi(t, y) H(t, y) dy = \int \Phi(t, y) D f'(U_t^{e_1}(dy)) + \int \frac{\partial \Phi(t, y)}{\partial t} H(t, y) dy. \quad (2.7)$$

Proof of Claim 2.1. Consider the limit of the incremental ratios. Integrate by parts in y before the limit, take then the limit in h and integrate by parts again in y . By the weak continuity of

$$t \mapsto D f'(U_t^{e_1}(dy))$$

one has

$$\int \Phi(t, y) D f'(U_t^{e_1}(dy)) = \lim_{h \downarrow 0} \frac{1}{h} \left\{ \iint \int_t^{t+h} \Phi(t, y) D_y f'(U(ds, dy)) \right\}. \quad (2.8a)$$

Remembering (2.3), (2.6) and then the definition (2.4b) of H one has

$$\begin{aligned}
& \iint_t^{t+h} \Phi(t, y) D_y f'(U(ds, dy)) = \\
& = \int \Phi(t, y) \left[\int_t^{t+h} f''(U(s, y)) \nu_y(ds) \vartheta(y) \right] dy \\
& = \int \Phi(t, y) [H(t+h, y) - H(t, y)] dy.
\end{aligned} \quad (2.8b)$$

Owing to (2.8) one can deduce that for \mathcal{L}^1 -a.e. t equation (2.1) holds:

$$\begin{aligned}
& \frac{d}{dt} \int \Phi(t, y) H(t, y) dy = \\
& = \lim_{h \rightarrow 0} \frac{1}{h} \left\{ \int \Phi(t+h, y) H(t+h, y) dy - \int \Phi(t, y) H(t, y) dy \right\} \\
& = \lim_{h \rightarrow 0} \frac{1}{h} \left\{ \int \Phi(t, y) [H(t+h, y) - H(t, y)] dy \right\} \\
& \quad + \int \lim_{h \rightarrow 0} \frac{\Phi(t+h, y) - \Phi(t, y)}{h} H(t+h, y) dy \\
& \stackrel{(2.8)}{=} \int \Phi(t, y) D f'(U_t^{e_1}(dy)) + \int \frac{\partial \Phi(t, y)}{\partial t} H(t, y) dy. \quad \square
\end{aligned}$$

The proof of the absolute continuity of suitable Lagrangian parameterizations is ended. \square

Proof of Lemma 22. We now prove that the PDE (1.1) holds in distributional sense. When $\partial_x u$ is a Radon measure, this implies by Vol'pert chain rule that u has locally bounded variation. For every test function $\varphi \in C_c^\infty(\Omega)$, one can apply in the integral

$$\langle \partial_t u + \partial_x f(u), \varphi \rangle = - \iint_\Omega \left[\frac{\partial \varphi(t, x)}{\partial t} u(t, x) + \frac{\partial \varphi(t, x)}{\partial x} f(u(t, x)) \right] dt dx$$

the following change of variables, that one can assume absolutely continuous by Corollary 21:

$$\Psi : \begin{pmatrix} t \\ y \end{pmatrix} \mapsto \begin{pmatrix} t \\ x \end{pmatrix} := \begin{pmatrix} t \\ \chi(t, y) \end{pmatrix}. \quad (2.9)$$

Denote $\Phi(t, y) = \varphi(i_{\chi(y)}(t))$ and $U(t, y) = u(i_{\chi(y)}(t))$. Remembering (2.5) one obtains

$$\begin{aligned}
\langle \partial_t u + \partial_x f(u), \varphi \rangle &= - \iint \frac{\partial \Phi(t, y)}{\partial t} U(t, y) D_y \chi(dt, dy) \\
& \quad + \iint f'(u(i_{\chi(y)}(t))) \frac{\partial \Phi(t, y)}{\partial y} U(t, y) dt dy \\
& \quad - \iint \frac{\partial \Phi(t, y)}{\partial y} f(U(t, y)) dt dy.
\end{aligned} \quad (2.10)$$

Step 1. y -derivatives. After integration by parts, the last two summands in the right-hand side of (2.10) are just

$$\iint \Phi(t, y) D_y [f(U) - f'(U)U] (dt, dy). \quad (2.11)$$

Notice that $f'(U)U$ is still a function with locally bounded variation, and that its derivative can be computed by Vol'pert chain rule: it is equal to

$$D_y [Uf'(U)] = [f'(U) + Uf''(U)] D_y U = D_y f(U) + U D_y f'(U),$$

and therefore (2.11) becomes

$$- \iint \Phi(t, y) U(t, y) D_y f'(U) (dt, dy). \quad (2.12)$$

Step 2. t -derivative. The first summand in the right-hand side of (2.10) is more complex and requires the properties of H in (2.4), (2.1). Notice that $\Phi(t, y)U(t, y)$ is absolutely continuous in time. By the additional regularity of H in (2.1) one has the integration by parts

$$\begin{aligned} & - \iint \frac{\partial \Phi(t, y)}{\partial t} U(t, y) D_y \chi(dt, dy) = \\ & = - \iint \frac{\partial \Phi(t, y)}{\partial t} U(t, y) H(t, y) dt dy \\ & = \iint \Phi(t, y) \frac{\partial U(t, y)}{\partial t} D_y \chi(dt, dy) + \\ & \quad + \iint \Phi(t, y) U(t, y) D_y f'(U) (dt, dy). \end{aligned} \quad (2.13)$$

Thanks to the absolute continuity of χ (that one can assume by Corollary 21), the term $\frac{\partial U(t, y)}{\partial t}$ in the third integral in (2.13) is just the Lagrangian source term $g_L = [g]_\chi$ evaluated at $i_{\chi(y)}(t)$. Therefore this integral can be rewritten just as

$$\iint \varphi(t, x) g(t, x) dt dx.$$

The fourth integral in (2.13) instead cancels the two remaining terms in (2.10), by their equivalent form (2.12). After this cancellation we find that (2.10) is just

$$\iint \varphi(t, x) [D_t u + D_x f(u)] (dt, dx) = \iint \varphi(t, x) g_L(t, x) dt dx. \quad \square$$

2.2. The case of continuous solutions: BV approximations

We provide in this section the proof that continuous Lagrangian solutions of the balance law (1.1) are also distributional solutions, without assuming any BV regularity. In order to prove it, we construct a sequence of approximations having bounded variation, so that we take advantage of §2.1. We omit here the correspondence of the source terms, discussed separately in [2].

Lemma 27. *Let u be a continuous Lagrangian solution of (1.1) with source term bounded by G . Then there exists a sequence of continuous functions $u_k(t, x)$, $k \in \mathbb{N}$, which are*

- functions of bounded variation;
- Lagrangian and Eulerian solutions of (1.1) with source terms bounded by G ;
- converging uniformly to u as $k \uparrow \infty$.

Corollary 28. *Let u be a continuous Lagrangian solution of (1.1). Then it is a continuous distributional solution of (1.1) and it does not dissipate entropy.*

The above corollary states in particular that u is the unique Kruřkov entropy solution to the Cauchy problem (when its distributional source term is assigned). We mention that in the case of the quadratic flux this statement can be derived by [7]: the authors provide a smooth approximation for which also the source term is converging in $\mathcal{L}^1(\mathbb{R}^2)$, refining a construction in [15] which extends to the Heisenberg group a technique originally introduced for the Euclidean setting by [9]. The construction we adopt here is more direct but rougher: sources do not converge.

Proof of Corollary 28. We exploit the approximation $\{u_k(t, x)\}_{k \in \mathbb{N}}$ given in Lemma 27. Consider any entropy-entropy flux pair $\eta, q \in C^1(\mathbb{R})$, that is, $q'(z) = \eta'(z)f'(z)$. Since each u_k is a function of bounded variation, Vol'pert chain rule yields

$$D_t \eta(u_k) + D_x(q(u_k)) = \eta'(u_k) (D_t u_k + D_x f(u_k)) = \eta'(u_k) g_{E_k},$$

where we set $g_{E_k} = D_t u_k + D_x f(u_k)$. Owing to Lemma 22, each g_{E_k} is given by a function which is bounded by the constant G in the assumption of the present corollary. Since u_k converges uniformly to u , one has

$$\eta'(u_k) g_{E_k} = \partial_t \eta(u_k) + \partial_x(q(u_k)) \rightarrow \partial_t \eta(u) + \partial_x(q(u)) \quad \text{in } \mathcal{D}'(\Omega).$$

Since the sequence $\{g_{E_k}\}_{k \in \mathbb{N}}$ is uniformly bounded, Banach–Alaoglu theorem implies that there exists a subsequence weak*-converging to some function $g_E \in L^\infty(\Omega)$, $\|g_E\|_{L^\infty(\Omega)} \leq G$: then necessarily

$$\partial_t \eta(u) + \partial_x(q(u)) = \eta'(u) g_E.$$

The Eulerian source g_E and the Lagrangian source g_L can be identified also in the limit under uniform convexity assumptions on the flux, see [2, §2.2]. Under the negligibility assumption on the inflection points of f (H), they are just compatible: see [2, §3 and §4.2]. \square

Proof of Lemma 27. We construct an approximation of u by a patching procedure. One needs first to construct the approximation on a patch, which is a strip delimited by two characteristics. In it, we require that at each fixed time the approximating function is monotone in x , and it coincides with u on the boundary of the strip. This allows to work with continuous functions having bounded variation. Repeating the construction in adjacent strips, when they get thinner the approximating functions converge to u uniformly.

We recall that u can be assumed compactly supported, see Notation 19.

We expose first the limiting procedure for constructing a monotone approximation within each strip, and then a second limiting procedure for converging to u when strips become thinner. In the first step we describe the second limiting procedure,

which is simpler, while from the second step on we describe how to provide the monotone approximations.

Step 1. Patches decomposition. Fix two characteristics $\chi_{y_1}^{e_1}(t)$, $\chi_{y_2}^{e_1}(t)$ and define the strip

$$S_{y_1 y_2} = \{(t, x) \in \mathbb{R}^+ \times \mathbb{R} : \chi(t, y_1) \leq x \leq \chi(t, y_2)\} \quad (2.14)$$

where $y_1 \leq y_2$. If one chooses for example $y_i = i\delta$ for $i \in \mathbb{Z}$ and some $\delta > 0$, then one has the decomposition

$$\mathbb{R}^+ \times \mathbb{R} = \cup_{i \in \mathbb{Z}} S_{y_i y_{i+1}}.$$

Let $|y_{i+1} - y_i| \leq \delta$. We construct in the next steps continuous functions u^δ which are

- (1) Lagrangian solutions, with a new Lagrangian source still bounded by G ;
- (2) equal to u on the curves $\chi_{y_i}^{e_1}(t)$, $i \in \mathbb{Z}$;
- (3) nondecreasing in the x -variable in each open t -section of the set

$$\left\{ (t, x) : \chi_{y_i}^{e_1}(t) < x < \chi_{y_{i+1}}^{e_1}(t), u(i_{\chi(y_i)}(t)) \leq u(i_{\chi(y_{i+1})}(t)) \right\};$$

- (4) nonincreasing in the x -variable in each open t -section of the set

$$\left\{ (t, x) : \chi_{y_i}^{e_1}(t) < x < \chi_{y_{i+1}}^{e_1}(t), u(i_{\chi(y_i)}(t)) \geq u(i_{\chi(y_{i+1})}(t)) \right\};$$

- (5) $\|u^\delta - u\|_{L^\infty} \leq \omega(\delta)$, where $\omega(\delta)$ is a δ -modulus of uniform continuity of

$$U(t, y) = u(i_{\chi_y^{e_1}}(t)).$$

From the monotonicity properties (3)-(4), if we apply the slicing theory for BV -functions we notice that the functions u^δ have locally bounded variation. Owing to Lemma 22 they are also Eulerian solutions with source terms which are uniformly bounded by G , the uniform bound for the Lagrangian sources owing to (1). By the uniform estimates (5) and the constrain (2) on the boundary of the strips, they converge uniformly to u as $\delta \downarrow 0$, proving the thesis.

Step 2. Monotone modification within a patch. We start the iterative procedure for constructing the approximations u^δ having bounded variation, that we describe within a fixed strip

$$S_{y_1 y_2} = \{(t, x) : \chi(t, y_1) < x < \chi(t, y_2)\}. \quad (2.15)$$

We modify in $S_{y_1 y_2}$ the given Lagrangian continuous solution in order to get a new continuous function which is still a Lagrangian solution, for a different source which is still bounded by G . The additional property that we are trying to get, piecewise, is a monotonicity in the x variable when t is fixed: we fix the values on the boundary curves $\chi_{y_1}^{e_1}(t)$, $\chi_{y_2}^{e_1}(t)$; inside the stripe $S_{y_1 y_2}$ we aim at substituting at each time $u_i^{e_1}(x)$ with a function $\tilde{u}_i^{e_1}(x)$ i) which is monotone in the x variable with values from $u(i_{\chi(y_1)}(t))$ to $u(i_{\chi(y_2)}(t))$ and ii) which is still a Lagrangian solution with source term bounded by G . This new function \tilde{u} is now defined inside the stripe $S_{y_1 y_2}$ with a limiting procedure pictured in Figure 2 below.

Let $\{(t_j, x_j)\}_{j \in \mathbb{N}}$ be a dense sequence of points within the strip $S_{y_1 y_2}$. The function \tilde{u} is defined within the strip $S_{y_1 y_2}$ as a uniform limit of functions \tilde{u}_j , for $j \in \mathbb{N}$, which we assign now recursively. We first state the basic operation that we will perform.




FIGURE 2. First steps of the iterative procedure: the bold line is the approximating function obtained by successive cuttings reaching monotonicity properties

Claim 29 (Basic cut). Suppose u is a Lagrangian solution of (1.1) with source term bounded by G and fix a characteristic curve $\bar{\gamma} = \chi(t, \bar{y})$, where χ is a Lagrangian parameterization of u . Then the two truncated functions, respectively from above and from below,

$$u_M(t, x) := u(t, x) \wedge u(i_{\bar{\gamma}}(t)), \quad u_m(t, x) := u(i_{\bar{\gamma}}(t)) \vee u(t, x)$$

are still Lagrangian solutions of (1.1) with source term bounded by G .

We postpone the proof of Claim 29 to Page 18. The basic cut allows the iterative procedure:

- set $m(t) := [u(i_{\chi(y_1)}(t))] \wedge [u(i_{\chi(y_2)}(t))]$;
- set $M(t) := [u(i_{\chi(y_1)}(t))] \vee [u(i_{\chi(y_2)}(t))]$;
- set $\tilde{u}_0(t, x) := u(t, x)$ for $(t, x) \in S_{y_1 y_2}$ and fix $\bar{\gamma}_0(t) := \chi(t, y_1(t))$;
- let $j \in \mathbb{N}$, set $v_j(t) = u(i_{\bar{\gamma}_{j-1}}(t))$, and define the truncated function

$$\tilde{u}_j(t, x) := \begin{cases} v_j(t) \vee \tilde{u}_{j-1}(t, x) \wedge M(t) & \text{if } u(i_{\chi(y_1)}(t)) = M(t) \\ & \text{and } \chi(t, y_1) \leq x \leq \bar{\gamma}_{j-1}(t), \\ m(t) \vee \tilde{u}_{j-1}(t, x) \wedge v_j(t) & \text{if } u(i_{\chi(y_1)}(t)) = m(t) \\ & \text{and } \chi(t, y_1) \leq x \leq \bar{\gamma}_{j-1}(t), \\ v_j(t) \vee \tilde{u}_{j-1}(t, x) \wedge M(t) & \text{if } u(i_{\chi(y_2)}(t)) = M(t) \\ & \text{and } \bar{\gamma}_{j-1}(t) \leq x \leq \chi(t, y_2), \\ m(t) \vee \tilde{u}_{j-1}(t, x) \wedge v_j(t) & \text{if } u(i_{\chi(y_1)}(t)) = m(t) \\ & \text{and } \bar{\gamma}_{j-1}(t) \leq x \leq \chi(t, y_2). \end{cases}$$

Basically, the strip $S_{y_1 y_2}$ is divided by $\bar{\gamma}_{j-1}$ into two sub-strips and in each sub-strip \tilde{u}_{j-1} is truncated from above and from below by the two boundary values on the sub-strips. Owing to Claim 29 the function \tilde{u}_j is a Lagrangian solution

of (1.1) with source term bounded by G . By the definition of Lagrangian solution one can fix $\tilde{\gamma}_j$ as

- a characteristic curve of \tilde{u}_j
- through the point (t_j, x_j)
- along which \tilde{u}_j is G -Lipschitz continuous and
- which does not cross the previously chosen characteristics $\chi(\cdot, y_1)$, $\tilde{\gamma}_1(t), \dots, \tilde{\gamma}_{j-1}(t)$, $\chi(\cdot, y_2)$; this means that any two characteristics of this set lie always on the same side of the pane with respect to each other, when they differ.

Note before proceeding that \tilde{u}_j is monotone at each fixed time on the $j + 2$ points

$$i_{\chi(y_1)}(t), \quad \{i_{\tilde{\gamma}_i(t)}\}_{i=1, \dots, j}, \quad i_{\chi(y_2)}(t). \quad (2.16)$$

Because of this monotonicity, at later steps of the iteration the values of the function \tilde{u}_j on the points (2.16) are not changed.

We obtained with the previous iterative procedure that

- each continuous function \tilde{u}_j is a Lagrangian solution and G still bounds its source, thanks to Claim 29;
- the whole sequence $\{\tilde{u}_j\}_{j \in \mathbb{N}}$ converges uniformly on $S_{y_1 y_2}$ to a function \tilde{u} , because u is uniformly continuous and the cutting procedure preserves the modulus of continuity: similarly to the next item, for $h > j$ one has the estimate

$$\|\tilde{u}_j - \tilde{u}_h\|_{L^\infty(S_{y_1 y_2})} \leq \omega \left(\text{diam} \left(S_{y_1 y_2} \setminus \text{Im} \left\{ i_{\chi(y_1)}, \{i_{\tilde{\gamma}_i}\}_{i=1, \dots, j}, i_{\chi(y_2)} \right\} \right) \right)$$

and the curves (2.16) become dense in $S_{y_1 y_2}$ by construction, so the right-hand side goes to 0;

- $\|u - \tilde{u}\|_{L^\infty(S_{y_1 y_2})} \leq \omega(y_2 - y_1)$, where ω is a modulus of uniform continuity of $U(t, y)$, because by construction we have

$$\begin{aligned} \|u - \tilde{u}\|_{L^\infty(S_{y_1 y_2})} &\leq \sup_{(t, x) \in S_{y_1 y_2}} \{|u(t, x) - u(i_{\chi(y_1)}(t))|, |u(t, x) - u(i_{\chi(y_2)}(t))|\} \\ &\leq \omega(y_2 - y_1) \leq \omega(\delta); \end{aligned} \quad (2.17)$$

- \tilde{u} is still a Lagrangian solution and that G still bounds its source by Corollary 47;
- \tilde{u} is monotone in the x -variable at each fixed t because each \tilde{u}_j is monotone on the points (2.16), which become dense in the interval $[\chi(t, y_1), \chi(t, y_1)]$. \square

We are finally left with the proof of Claim 29.

Proof of Claim 29. Owing to Lemma 17, $u_m(t, x) := u(i_{\tilde{\gamma}}(t)) \vee u(t, x)$ is a Lagrangian solution provided that we exhibit a characteristic curve, along which u_m is G -Lipschitz continuous, through any point (t, x) . For simplifying the exposition suppose that the set $\{f'(u_m) < f'(u)\}$ is empty, if not this region is treated similarly

to below as a second step. Set

$$\lambda_m(t, x) := f'(u_m(t, x)) \geq \lambda(t, x) := f'(u(t, x)).$$

Consider now the set Γ of curves through a point (\bar{t}, \bar{x}) obtained as the closure in the C^1 norm of the set all curves $\gamma \in C^1(\mathbb{R}^+)$ such that

- $\gamma(\bar{t}) = \bar{x}$;
- there exist an integer j , intervals $I_1, \dots, I_j, J_0, \dots, J_j$ and values $c_1, \dots, c_j, z_0, \dots, z_j$ such that $J_0 \cup I_1 \cup J_1 \cup \dots \cup J_j = \mathbb{R}^+$ and

$$\begin{cases} \gamma(t) = \chi(t, \tilde{\gamma}(t)) + c_i, & \lambda_m(i_{\tilde{\gamma}}(t)) > \lambda(i_{\tilde{\gamma}}(t)) & \text{if } t \in I_i, i = 1, \dots, j, \\ \gamma(t) = \chi(t, z_i) & & \text{if } t \in J_i, i = 0, \dots, j. \end{cases}$$

This set of curves is not empty, for example because the curves of the Lagrangian parameterization χ through (t, x) belongs to it. Moreover, the function u_m is G -Lipschitz continuous on each γ described within the brackets: indeed, where $\lambda_m > \lambda$ necessarily $u_m \neq u$ and hence by definition of u_m

$$u_m(\gamma(t)) = \begin{cases} u(i_{\tilde{\gamma}}(t)) & \text{if } t \in I_1 \cup \dots \cup I_j, \\ u(i_{\chi(z_i)}(t)) & \text{if } t \in J_i, i = 0, \dots, j, \end{cases}$$

and each function $u(i_{\tilde{\gamma}}(t)), u(i_{\chi(z_0)}(t)), \dots, u(i_{\chi(z_j)}(t))$ is G -Lipschitz continuous by assumption. As a consequence, u_m is G -Lipschitz continuous on each element of the closure Γ . The curves defined by

$$\begin{aligned} \gamma_m^+(t) &:= \max_{\gamma \in \Gamma} \gamma(t) & \text{for } t \geq \bar{t}, \\ \gamma_m^-(t) &:= \min_{\gamma \in \Gamma} \gamma(t) & \text{for } t < \bar{t}, \end{aligned}$$

still belongs to Γ , if suitably prolonged. In particular, u_m is G -Lipschitz continuous along γ_m . This concludes the proof observing that γ_m^+ is necessarily a forward characteristic curve of u_m , and γ_m^- a backward one, through (\bar{t}, \bar{x}) . Indeed, each γ in the definition of Γ is a C^1 curve whose slope satisfies the implication

$$\dot{\gamma} \in \{\lambda \circ i_{\tilde{\gamma}}, \lambda_m \circ i_{\tilde{\gamma}}\} \Rightarrow \dot{\gamma}_m^+, \dot{\gamma}_m^- \in \{\lambda \circ i_{\tilde{\gamma}}, \lambda_m \circ i_{\tilde{\gamma}}\}.$$

If we had $\dot{\gamma}_m^+(t') = \lambda \circ i_{\tilde{\gamma}_m^+}(t') < \lambda_m \circ i_{\tilde{\gamma}_m^+}(t')$ at some time $t' > \bar{t}$, then we would contradict the extremity in the definition of γ_m^+ : considering

$$s' = \bar{t} \vee \max\{s < t' : \dot{\gamma}_m^+(s) = \lambda_m \circ i_{\tilde{\gamma}_m^+}(s)\},$$

one can verify that there is an element of Γ which satisfies

$$\tilde{\gamma}_m^+(t) = \begin{cases} \gamma_m^+(t) & \text{for } \bar{t} \leq t \leq s', \\ \tilde{\gamma}_\tau(t) := \tilde{\gamma}(t) - [\tilde{\gamma}(s') - \gamma_m^+(s')] & \text{for } s' < t \leq s'', \end{cases}$$

where

$$s'' := t' \wedge \min\{s > s' : \lambda_m \circ i_{\tilde{\gamma}_\tau}(s) \neq \lambda \circ i_{\tilde{\gamma}_\tau}(s)\}.$$

Notice that $\tilde{\gamma}_m^+$ is bigger than γ_m at time s'' because $\dot{\tilde{\gamma}}_m^+(s'') = \dot{\gamma}_m^+(s'')$ and by construction

$$\dot{\tilde{\gamma}}_m^+ = \lambda_m \circ i_{\tilde{\gamma}} = \lambda_m \circ i_{\tilde{\gamma}_m^+}(s) > \lambda \circ i_{\tilde{\gamma}_m^+}(s) = \dot{\gamma}_m^+ \quad \text{for } s' < t < s''.$$

The other possibilities contradict analogously the definition of γ_m . \square

3. DISTRIBUTIONAL SOLUTIONS ARE BROAD SOLUTIONS, IF INFLECTIONS ARE NEGLIGIBLE

We provide in this section regularity results holding under the assumption that f has negligible inflection points: we prove that u is Lipschitz continuous along every characteristic curve and there exists a universal source which is fine for every Lagrangian parameterization one chooses.

Without assumptions on inflection points, later §4 shows that distributional solutions of

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g_E(t, x) \quad (1.1)$$

(with $f \in C^2(\mathbb{R})$, $|g_E(t, x)| \leq G$) are also Lagrangian solutions. Being a Lagrangian solution allows to study u with tools from ODEs, but it is not completely satisfactory by itself because one should be a priori careful in choosing the right Lagrangian parameterization, and the correct source related to the parameterization: the results of the present section are richer because here *any* Lagrangian parameterization is allowed.

Being local arguments, we simplify the setting by assuming $\Omega = \mathbb{R}^+ \times \mathbb{R}$, u compactly supported.

3.1. Lipschitz regularity along characteristics

In the present section we point out that u is Lipschitz continuous along characteristic curves if inflection points of f are negligible:

$$\mathcal{L}^1(\text{clos}(\text{Infl}(f))) = 0. \quad (\text{H})$$

See Example [2, §4.2] for an example when (H) fails.

Theorem 30. *Assume that the non-vanishing condition (H) holds. Then any continuous distributional solution u of (1.1) is G -Lipschitz continuous along any characteristic curve of u .*

Proof. It takes a while to realize that the following is a partition of the real line into the regions

$$D^+ := \text{r.i.} \left(\{z \in \mathbb{R} : \exists \bar{h} > 0 \text{ s.t. } f(z+h) - f(z) \geq f'(z)h \forall h \in [-\bar{h}, \bar{h}]\} \right),$$

$$D^- := \text{r.i.} \left(\{z \in \mathbb{R} : \exists \bar{h} > 0 \text{ s.t. } f(z+h) - f(z) \leq f'(z)h \forall h \in [-\bar{h}, \bar{h}]\} \right),$$

$$N := \mathbb{R} \setminus (D^+ \cup D^-) = \text{clos}(\text{Infl}(f)).$$

By assumption N is Lebesgue negligible.

Consider any characteristic curve $i_\gamma(t) = (t, \gamma(t))$ where $\dot{\gamma}(t) = f'(u(t, \gamma(t)))$ and $t \in \mathbb{R}^+$. We first follow a similar computation in [8] which shows that $u(t, \gamma(t))$ is $\|g\|_\infty$ -Lipschitz continuous on the connected components of the open set $(u \circ i_\gamma)^{-1}(D^+)$, as in [5].

Focus on the domain bounded by the curves $i_\gamma(t)$, $i_{\gamma+\varepsilon}(t)$ between times $\sigma < \tau$. By integrating suitable test functions converging to the indicator of the region described




FIGURE 3. Balances on characteristic regions

in Figure 3.1 we obtain the equality

$$\begin{aligned} & \int_{\gamma(\tau)}^{\gamma(\tau)+\varepsilon} u(\tau, x) dx - \int_{\gamma(\sigma)}^{\gamma(\sigma)+\varepsilon} u(\sigma, x) dx - \int_{\sigma}^{\tau} \int_{\gamma(t)}^{\gamma(t)+\varepsilon} g(t, x) dx dt \\ &= - \int_{\sigma}^{\tau} [f(u(i_{\gamma+\varepsilon}(t))) - f(u(i_{\gamma}(t)))] \\ & \quad - f'(u(i_{\gamma}(t))) [u(i_{\gamma+\varepsilon}(t)) - u(i_{\gamma}(t))] dt \\ &= - \int_{\sigma}^{\tau} f''(\xi) \frac{[u(i_{\gamma+\varepsilon}(t)) - u(i_{\gamma}(t))]^2}{2} dt, \end{aligned}$$

where in the last integral $\xi = \xi(t) \in [u(i_{\gamma}(t)) - u(i_{\gamma+\varepsilon}(t))]$.

If either $u(\sigma, \gamma(\sigma))$ or $u(\tau, \gamma(\tau))$ belong to D^+ , by definition of D^+ for $\tau - \sigma$ sufficiently small the right-hand side is nonpositive: we obtain thus the inequality

$$\begin{aligned} & \int_{\gamma(\tau)}^{\gamma(\tau)+\varepsilon} u(\tau, x) dx - \int_{\gamma(\sigma)}^{\gamma(\sigma)+\varepsilon} u(\sigma, x) dx \\ & \leq \int_{\sigma}^{\tau} \int_{\gamma(t)}^{\gamma(t)+\varepsilon} g(t, x) dx dt \leq \varepsilon \|g\|_\infty |\tau - \sigma|. \end{aligned}$$

Dividing by ε , and taking the limit as $\varepsilon \downarrow 0$, the assumption that u is continuous yields

$$u(\tau, \gamma(\tau)) - u(\sigma, \gamma(\sigma)) \leq \|g\|_\infty |\tau - \sigma|.$$

The converse inequality is obtained by considering the similar region between $i_{\gamma-\varepsilon}(t)$, $i_\gamma(t)$: indeed this leads to an equation analogous to (3.1), but with right-hand side having opposite sign.

We conclude from the above analysis that $u(i_\gamma(t))$ is $\|g\|_\infty$ -Lipschitz continuous in a neighborhood of any point belonging to the inverse image of D^+ . The same holds in an analogous way for D^- . This local Lipschitz continuity can be equivalently stated by the inequality

$$\mathcal{L}^1(u(i_\gamma(B))) \leq G \mathcal{L}^1(B) \quad (3.2)$$

for all Borel subsets B of the open set $O = (u \circ i_\gamma)^{-1}(D^+ \cup D^-)$. The thesis finally follows by the negligibility of N : for every $t_1 \leq t_2$

$$\begin{aligned} |u(i_\gamma(t_2)) - u(i_\gamma(t_1))| &\leq \mathcal{L}^1(u(i_\gamma([t_1, t_2]))) \\ &= \mathcal{L}^1(u(i_\gamma([t_1, t_2] \cap O))) + \mathcal{L}^1(N) \\ &\stackrel{(3.2)}{\leq} G \mathcal{L}^1([t_1, t_2] \cap O) \leq G(t_2 - t_1). \quad \square \end{aligned}$$

Remark 31. If u is 1/2-Hölder continuous, we see from (3.1) that u is Lipschitz along characteristics independently of any assumptions on inflection points of f .

3.2. Construction of a universal source

We now assume the negligibility of inflection points (H). Under this assumption, we generalize [5, §6] and we construct for general fluxes satisfying (H) a source term for the broad formulation, without discussing its compatibility with the distributional source. Namely, we show that there exists $\mathbf{g} \in \mathcal{L}^\infty(\Omega)$ such that

$$\frac{d}{dt}u(i_\gamma(t)) = \mathbf{g}(i_\gamma(t)) \quad \text{in } \mathcal{D}(i_\gamma^{-1}(\Omega)) \text{ for every characteristic curve } \gamma. \quad (3.3a)$$

The compatibility of the sources will be instead matter of [2, §3], where we prove that when inflection points are negligible there is a choice of such \mathbf{g} so that moreover

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = \mathbf{g}(t, x) \quad \text{in } \mathcal{D}(\Omega). \quad (3.3b)$$

We deal here in §3.2 only with the ODE property (3.3a). We call such $\mathbf{g}_B = [\mathbf{g}]_\lambda$ universal source term, and (3.3a) shows that u is a broad solution of (1.1). We mention nevertheless that if f is not α -convex, $\alpha > 1$, then (3.3a) does not identify in general a distribution, because there can be an \mathcal{L}^2 -positive measure set of points where u is not differentiable along characteristics: in this set the proper definition of the source will come from (3.3b).

Two remarks before starting. Owing to §3.1, under the sharp vanishing condition (H) on inflection points of f one gains G -Lipschitz continuity along characteristic curves for any continuous distributional solution u to the balance law

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g_E(t, x). \quad (1.1)$$

It is not of course possible to require that the reduction of the balance law on characteristics is satisfied for every $\mathbf{g} \in \mathcal{L}^\infty(\mathbb{R}^+ \times \mathbb{R})$ such that $g_E = [\mathbf{g}]$, because altering $\mathbf{g}(t, x)$ on a curve provides the same distribution $[\mathbf{g}]$: this is why we need to select a good representative. Without the negligibility (H) the source term of a Lagrangian parameterization might not work with a different Lagrangian parameterization and there may exist no broad solution, see [2, §4.3].

We assume therefore the negligibility of inflection points (H) and we proceed as follows: in §3.2.1 we construct a *Souslin* function \mathbf{g} , which intuitively must satisfy (3.3a); in §3.2.2 we construct an analogous *Borel* function $\hat{\mathbf{g}}$, which is stronger but more technical; in §3.2.3 we prove that the functions \mathbf{g} and $\hat{\mathbf{g}}$ do satisfy (3.3a).

The construction for the compatibility condition (3.3b) comes in [2, §3].

3.2.1. Souslin selection

This is the first idea: to define pointwise, but in a measurable way, a function $\hat{g}(t, x)$ such that t is a Lebesgue point for the derivative of the composition $t \mapsto u(t, \gamma(t))$, with γ a characteristic function through (t, x) , whenever there exists one satisfying this differentiability property. As we just consider the derivative of this composition at t fixed, we focus on the curve only in a neighborhood of t and, for notational convenience, we translate its domain to a neighborhood of the origin. Therefore, fixed some $\delta > 0$, one applies a selection theorem to the set

$$\mathcal{G} \subset [\delta, +\infty) \times \mathbb{R} \times C^1([-\delta, \delta]; \mathbb{R}) \times [-G, G], \quad \mathcal{G} := \mathcal{C} \cap \mathcal{D}$$

where

$$\mathcal{C} := \left\{ (t, x, \gamma, \zeta) : \gamma(0) = x, \dot{\gamma}(s) = \lambda(t + s, \gamma(s)) \right\} \quad (3.4a)$$

$$\mathcal{D} := \left\{ (t, x, \gamma, \zeta) : \zeta = \lim_{\sigma \downarrow 0} \frac{u(t \pm \sigma, \gamma(\pm \sigma)) - u(t, x)}{\pm \sigma} \right\}. \quad (3.4b)$$

We first need a technical but important lemma about \mathcal{G} . The selection theorem will follow.

Lemma 32. \mathcal{G} is Borel.

Proof. Focus first on the components (t, x, γ) . The set \mathcal{C} is closed thanks to the continuity of λ .

We discretize the limit in the variable ζ , so that \mathcal{D} is described as an $F_{\sigma\delta}$ -set.

Claim 33. Existence and the values of the following two limits are the same:

$$\lim_{h \downarrow 0} \frac{u(t+h, \gamma(h)) - u(t, x)}{h}, \quad (3.5)$$

$$\lim_{n \rightarrow \infty} \frac{u(t+h_n, \gamma(h_n)) - u(t, x)}{h_n}, \quad h_{n+1} := h_n - h_n^2, \quad h_1 := 1/2. \quad (3.6)$$

One can similarly have the full limit for $h \rightarrow 0$ instead of $h \downarrow 0$, that we study for simplicity.

Proof of Claim 33. By Theorem 30, u is $\|g\|_\infty$ -Lipschitz continuous on characteristic curves. Setting $(t, x) = (0, 0)$ for notational convenience, then for every $h \in (h_{n+1}, h_n]$ there holds

$$\begin{aligned} &\left| \frac{u(h, \gamma(h)) - u(0, 0)}{h} - \frac{u(h_n, \gamma(h_n)) - u(0, 0)}{h_n} \right| \\ &= \left| \left(\frac{1}{h} - \frac{1}{h_n} \right) [u(h, \gamma(h)) - u(0, 0)] - \frac{1}{h_n} [u(h_n, \gamma(h_n)) - u(h, \gamma(h))] \right| \\ &\leq 2G \frac{h_n - h}{h_n}. \end{aligned}$$

By construction however

$$|h_n - h| \leq |h_n - h_{n+1}| = h_n^2,$$

yielding that the existence of the limit along $\{h_n\}_n$ implies the existence of the limit for any $h \downarrow 0$. \square

Notice that the claim would not hold choosing a generic $\tilde{h}_n \downarrow 0$ instead of $\{h_n\}_n$. After observing that the limit is discrete, the classical differentiability constraint in (3.4b) is

$$\forall k \exists n \text{ s.t. } \forall \bar{n} \geq n \quad \left| \zeta - \frac{u(t \pm h_{\bar{n}}, \gamma(h_{\bar{n}})) - u(t, x)}{\pm h_{\bar{n}}} \right| \leq 2^{-k}.$$

Therefore, \mathcal{D} is equivalently described as the following $F_{\sigma\delta}$ set

$$\mathcal{D} = \bigcap_{k \in \mathbb{N}} \bigcup_{n \in \mathbb{N}} \bigcap_{\bar{n} \geq n} \left\{ (t, x, \gamma, \zeta) : \left| \zeta - \frac{u(t \pm h_{\bar{n}}, \gamma(h_{\bar{n}})) - u(t, x)}{\pm h_{\bar{n}}} \right| \leq 2^{-k} \right\}.$$

Since the set within brackets is closed, $\mathcal{G} = \mathcal{C} \cap \mathcal{D}$ is Borel. \square

Let E be the projection of \mathcal{G} on the first two components $\mathbb{R}^+ \times \mathbb{R}$. The set E is the set of points where there exists an absolutely continuous (time-translated) characteristic curve having 0 as a density point for the derivative of $u(t+s, \gamma_{(t,x)}(s))$. The selection theorem below assigns to every point (t, x) where possible, which is to every point in E , an absolutely continuous integral curve $\gamma_{(t,x)}(s)$ for the ODE

$$\dot{\gamma}_{(t,x)}(s) = \lambda(t + s, \gamma_{(t,x)}(s))$$

together with the Souslin function

$$(t, x) \mapsto \mathbf{g}(t, x) = \frac{d}{ds} u(t + s, \gamma_{(t,x)}(s)) \Big|_{s=0}. \quad (3.7)$$

Remark 34. We comment on what information on E comes from hypothesis on f :

(1) If f is α -convex, we will observe in [2, §3.2] that the projection E of \mathcal{G} on $\mathbb{R}^+ \times \mathbb{R}$ has full measure. This follows for the case of quadratic flux by a Rademacher theorem in the context of the Heisenberg group [11, 5].

(2) If f is even strictly but not uniformly convex [2, §4.2] shows that E may fail to have full measure. If (the closure of) inflection points of f are negligible, however, the Lipschitz continuity of u along characteristics of Theorem 30 implies that

$$\mathcal{H}^1(i_\gamma(\mathbb{R}) \setminus E) = 0$$

for every characteristic curve $\gamma(t)$.

(3) For general fluxes not only E may not have full \mathcal{L}^2 -measure, but also $i_\gamma(\mathbb{R}) \setminus E$ may not have full \mathcal{H}^1 -measure for some characteristic curve γ along which u is not Lipschitz-continuous, see [2, §4.3].

The set E is considered also in [2, §3] for the compatibility of the source terms.

Corollary 35 (Selection theorem). *For every $\delta > 0$, there exists a map*

$$(t, x) \mapsto (\gamma_{(t,x)}(s), \mathbf{g}(t, x)) \in C^1([-\delta, \delta]; \mathbb{R}) \times [-G, G]$$

where

$$(t, x) \in E \subset [\delta, +\infty) \times \mathbb{R}, \quad \gamma_{(t,x)} \in C^1([-\delta, \delta]; \mathbb{R}), \quad \mathbf{g}(t, x) \in [-G, G],$$

which is measurable for the σ -algebra generated by analytic sets and which satisfies by definition

$$(t, x, \gamma_{(t,x)}(s), \mathbf{g}(t, x)) \in \mathcal{G}.$$

Proof. The Borel measurability of \mathcal{G} proved in Lemma 32 allows to apply to \mathcal{G} von Neumann selection theorem [18, Theorem 5.5.2], from [16], which provides the thesis. \square

Definition 36. We define as a Souslin universal source the function

$$\mathbf{g}(t, x) = \mathbf{g}_{t,x} = \begin{cases} 0 & (t, x) \notin E \\ 0 & u(t, x) \in N \\ \mathbf{g}_{t,x} & (t, x) \in E, u(t, x) \notin N. \end{cases}$$

The importance of the above selection theorem is due to the following relation.

Theorem 37. *Assume that $\mathcal{L}^1(\text{clos}(\text{Infl}(f))) = 0$. Then for every absolutely continuous integral curve γ of the ODE $\dot{\gamma} = \lambda(i_\gamma)$, one has that $\mathbf{g}(i_\gamma)$ is well defined \mathcal{L}^1 -a.e. and satisfies*

$$u(i_\gamma(s)) - u(i_\gamma(r)) = \int_r^s \mathbf{g}(i_\gamma(t)) dt \quad \forall 0 \leq r \leq s.$$

Theorem 37 is fairly nontrivial because in (3.7) the universal source $\mathbf{g}(t, x)$ is defined as the derivative of u along a chosen curve $\gamma_{(t,x)}$ which changes changing the point (t, x) , and it is not even defined on a full measure set! What is relevant for the theorem is that the set where \mathbf{g} is not defined, or not uniquely defined, is negligible along any characteristic curve, which is that

$$\mathbf{g}_B = [\mathbf{g}]_\lambda$$

is well defined independently of the selection we have made. Different selections may change \mathbf{g} , but not \mathbf{g}_B . We postpone the proof of the theorem and of this fact to §3.2.3, after showing that it is possible to define a Borel selection $\hat{\mathbf{g}}$. Theorem 37 implies that \mathbf{g} and $\hat{\mathbf{g}}$ give the same \mathbf{g}_B .

3.2.2. Borel selection

Before proving Theorem 37, for the sake of completeness we show that one can define as well a Borel function, that we denote by $\hat{\mathbf{g}}(t, x) = \hat{\mathbf{g}}_{t,x}$, for which Theorem 37 still holds. This requires a bit more work than the previous argument: we do not associate immediately to each point (where it is possible) an eligible curve and the derivative of u along it, but something which must be close to it. We find then with the proof of Theorem 37 that we end up with the same class $\mathbf{g}_B = [\hat{\mathbf{g}}]_\lambda$.

Lemma 38. *The (t, x) -projection E of \mathcal{G} is Borel. For every $\varepsilon > 0$ there exists a Borel function*

$$(t, x) \mapsto (\gamma_{\varepsilon,t,x}, \mathbf{g}_{\varepsilon,t,x})$$

with

$$(t, x) \in S, \quad \gamma_{\varepsilon,t,x} \in C^1([-\delta, \delta]), \quad \mathbf{g}_{\varepsilon,t,x} \in [-G - \varepsilon, G + \varepsilon]$$

such that $(t, x, \gamma_{\varepsilon, t, x}, \mathfrak{g}_{\varepsilon, t, x}) \in \mathcal{C}$ of (3.4a) and such that for $|h|$ sufficiently small

$$\left| \mathfrak{g}_{\varepsilon, t, x} - \frac{u(t \pm h, \gamma_{\varepsilon, t, x}(\pm h)) - u(t, x)}{\pm h} \right| < \varepsilon.$$

Definition 39. We define as a Borel universal source the function

$$\hat{\mathfrak{g}}(t, x) = \hat{\mathfrak{g}}_{t, x} = \begin{cases} 0 & \text{if } (t, x) \notin E \text{ or } u(t, x) \in N, \\ \liminf_{\varepsilon \downarrow 0} \mathfrak{g}_{\varepsilon, t, x} & \text{if } (t, x) \in E \text{ and } u(t, x) \notin N, \end{cases}$$

where $\mathfrak{g}_{\varepsilon, t, x}$ is fixed in Lemma 38.

Proof of Lemma 38. We remind the following selection theorem [18, Th. 5.12.1].

Theorem 40 (Arsenin–Kunugui). *Let $B \subset X \times Y$ be a Borel set, X, Y Polish, such that B_x is σ -compact for every x . Then the projection on X of B is Borel, and B admits a Borel function $s : P_X B \rightarrow Y$ such that $(x, s(x)) \in B$ for all x in the projection $P_X B$.*

We verify the hypothesis of, and we apply the above selection theorem to the set

$$\bigcup_{n \in \mathbb{N}} \bigcap_{m > n} \left\{ (t, x, \gamma, \zeta) \in \mathcal{C} : \left| \zeta - \frac{u(t \pm h_m, \gamma(\pm h_m)) - u(t, x)}{\pm h_m} \right| \leq \frac{\varepsilon}{2} \right\}, \quad (3.8)$$

where \mathcal{C} was defined in (3.4a) and $\{h_n\}_{n \in \mathbb{N}}$ immediately below that in (3.6). The section

$$\mathcal{C}_{(\bar{t}, \bar{x})} = \{(\bar{t}, \bar{x}, \gamma, \zeta) : \gamma(0) = \bar{x}, \dot{\gamma}(s) = \lambda(t + s, \gamma(s))\}$$

is locally compact as a consequence of the Arzelà–Ascoli theorem, because by the boundedness of λ the curves are equi-bounded and equi-Lipschitz continuous, and by the continuity of λ when they converge uniformly they also converge in $C^1([-\delta, \delta])$. For t, h fixed the set

$$\left\{ (\gamma, \zeta) : \left| \zeta - \frac{u(t \pm h, \gamma(\pm h)) - u(t, x)}{\pm h} \right| \leq \frac{\varepsilon}{2} \right\}$$

is closed, therefore its intersection with $\mathcal{C}_{(\bar{t}, \bar{x})}$ is compact: this proves that each (t, x) -section of (3.8) is σ -compact. The hypothesis of the theorem are satisfied: it provides that the projection E of (3.8) on the first factor $\mathbb{R}^+ \times \mathbb{R} \ni (t, x)$ is Borel and that there exists a Borel subset of (3.8) which is the graph of a function s defined on E . Since the projection from that graph to the first components $\mathbb{R}^+ \times \mathbb{R}$ is one-to-one and continuous, the function s is a Borel section of (3.8), concluding our statement. \square

3.2.3. Proof of Theorem 37

We provide here the proof of Theorem 37 with $\mathfrak{g}_{t, x}$ either the Borel or the Souslin one. Let us introduce the notation. We consider:

- $\bar{\gamma}(s)$ a characteristic curve for the balance law through $(t, x) = (t, \bar{\gamma}(t))$;
- $\bar{\zeta}(s) = \frac{d}{ds} u(s, \bar{\gamma}(s))$ the derivative of u along $\bar{\gamma}$, where it exists;

- either $\gamma_{\varepsilon, t, x}(s)$ or $\gamma_{t, x}(s)$: the characteristic curve of u through (t, x) given either by Lemma 38 or by Corollary 35 (fix for example $\gamma_{\varepsilon, t, x}(s)$, which is more complex);
- $\zeta_{\varepsilon}(t + s) = \frac{d}{ds} u(t + s, \gamma_{\varepsilon, t, x}(s))$ the derivative of u along $\gamma_{\varepsilon, t, x}$, where it exists; where the derivative does not exist, set the function equal to e.g. 0;
- either $\hat{\mathfrak{g}}(t, x)$ of Definition 39 or $\mathfrak{g}(t, x)$ of Corollary 35 (fix $\hat{\mathfrak{g}}(t, x)$, as we are showing the proof with the Borel selection of §3.2.2).

We indeed know from Theorem 30 that $u(i_{\bar{\gamma}}(s))$ and $u(i_{\gamma_{\varepsilon, t, x}}(s))$ are G -Lipschitz continuous. We prove first that for almost every t the derivative of $u(i_{\bar{\gamma}}(t))$ is precisely $\hat{\mathfrak{g}}(i_{\bar{\gamma}}(t))$ if $u(i_{\bar{\gamma}}(t))$ is not an inflection point of f . After that, we exploit again the negligibility assumption $\mathcal{L}^1(\text{clos}(\text{Infl}(f))) = 0$ on the inflection points of f and we conclude

$$u(i_{\bar{\gamma}}(s)) - u(i_{\bar{\gamma}}(r)) = \int_r^s \hat{\mathfrak{g}}(i_{\bar{\gamma}}(t)) dt \quad \forall 0 \leq r \leq s.$$

Step 1: Countable decomposition. We give a countable covering of the set of Lebesgue points t where the derivative $\bar{\zeta}(t)$ of $u(i_{\bar{\gamma}}(t))$ exists but differs from $\hat{\mathfrak{g}}_{i_{\bar{\gamma}}(t)}$. The set can be described as

$$\bigcup_{\varepsilon > 0} \left\{ t : \exists \bar{\zeta}(t) = \lim_{\sigma \rightarrow 0} \frac{u(i_{\bar{\gamma}}(t + \sigma)) - u(i_{\bar{\gamma}}(t))}{\sigma}, \left| \hat{\mathfrak{g}}_{i_{\bar{\gamma}}(t)} - \bar{\zeta}(t) \right| \geq \varepsilon \right\}.$$

In particular, dropping the condition that the derivative $\bar{\zeta}(t)$ of $u(i_{\bar{\gamma}}(t))$ exists at t we notice that this set is contained in

$$\bigcup_{\varepsilon > 0} \bigcup_{n \in \mathbb{N}} \left\{ t : \left| \hat{\mathfrak{g}}_{i_{\bar{\gamma}}(t)} - \frac{1}{\sigma} \int_t^{t+\sigma} \bar{\zeta} \right| \geq \varepsilon, \left| \hat{\mathfrak{g}}_{i_{\bar{\gamma}}(t)} - \frac{1}{\sigma} \int_{t-\sigma}^t \bar{\zeta} \right| \geq \varepsilon \forall \sigma \in (0, 2^{-n}) \right\}.$$

The proof now needs a further index because we are working with the Borel selection. If we recall the Definition 39 of $\hat{\mathfrak{g}}_{t, x}$, and observe that along the characteristic curve $\gamma_{\varepsilon, t, \bar{\gamma}(t)}$ there holds

$$u(i_{\gamma_{\varepsilon, t, x}}(t \pm \sigma)) - u(i_{\gamma_{\varepsilon, t, x}}(t)) = \int_t^{t \pm \sigma} \zeta_{\varepsilon},$$

then we can add a condition which is always satisfied, and the last union can be rewritten as

$$\bigcup_{\tilde{\varepsilon} < \varepsilon} \bigcup_{n \in \mathbb{N}} \left\{ t : \left| \hat{\mathfrak{g}}_{i_{\bar{\gamma}}(t)} - \frac{1}{\pm \sigma} \int_t^{t \pm \sigma} \bar{\zeta} \right| \geq 3\varepsilon, \left| \mathfrak{g}_{\tilde{\varepsilon}, i_{\bar{\gamma}}(t)} - \frac{1}{\pm \sigma} \int_t^{t \pm \sigma} \zeta_{\tilde{\varepsilon}} \right| < \varepsilon \quad \forall \sigma \in (0, 2^{-n}) \right\}.$$

The union can as well be done on any sequences $\tilde{\varepsilon}_k < \varepsilon_k$ with $\varepsilon_k \downarrow 0$: if

$$|\hat{\mathfrak{g}}_{t, x} - \mathfrak{g}_{\tilde{\varepsilon}_k, t, x}| \leq \varepsilon_k / 3$$

then one has the equivalent expression

$$\bigcup_{\tilde{\varepsilon}_k < \varepsilon_k} \bigcup_{n \in \mathbb{N}} \left\{ t : \left| \hat{\mathbf{g}}_{i_{\bar{\gamma}}}(t) - \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \bar{\zeta} \right| \geq 3\varepsilon_k, \right. \\ \left. \left| \hat{\mathbf{g}}_{i_{\bar{\gamma}}}(t) - \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \zeta_{\tilde{\varepsilon}_k} \right| < \varepsilon_k \quad \forall \sigma \in (0, 2^{-n}) \right\}.$$

We arrived to the countable covering that we wanted to prove in this step. If one is considering the Souslin selection clearly $\gamma_{\tilde{\varepsilon}, t, x} = \gamma_{t, x}$ and $\mathbf{g}_{\tilde{\varepsilon}, t, \bar{\gamma}(t)} = \mathbf{g}_{t, \bar{\gamma}(t)} = \zeta_{\tilde{\varepsilon}_k}(t)$.

Step 2: Reduction argument. We prove that the set

$$\left\{ t : \hat{\mathbf{g}}(i_{\bar{\gamma}}(t)) > \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \bar{\zeta} + 3\varepsilon, \right. \\ \left. \left| \hat{\mathbf{g}}(i_{\bar{\gamma}}(t)) - \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \zeta_{\tilde{\varepsilon}} \right| < \varepsilon \quad \forall \sigma \in (0, 2^{-n}) \right\} \quad (3.9)$$

cannot contain two points t_1, t_2 with $|t_1 - t_2| \leq 2^{-n}$. The case

$$\hat{\mathbf{g}}(i_{\bar{\gamma}}(t)) < \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \bar{\zeta} - 3\varepsilon$$

is similar, arguing backwards in time. Then the thesis will follow: by the previous step, the set of times where the derivative of $u(i_{\bar{\gamma}}(t))$ exists and it is different from $\hat{\mathbf{g}}_{i_{\bar{\gamma}}(t)}$ will be at most countable. Therefore the derivative of $u(i_{\bar{\gamma}}(t))$ will be almost everywhere precisely $\hat{\mathbf{g}}_{i_{\bar{\gamma}}(t)}$.

Step 3: Analysis of the single sets. By contradiction, assume that (3.9) contains two such points, for example $t_1 = 0, t_2 = \rho$. Then, essentially two cases may occur.

Substep 3.1: Concavity/convexity region. We first consider the open region where $f''(u) \geq 0$. (The open region $f''(u) \leq 0$ is entirely similar.) The restriction to the open set is allowed because the argument is local: we consider later also the region of inflection points. In particular, in this step we consider $f'(u)$ monotone in u , in particular nondecreasing.

Compare $\bar{\gamma}$ with the two curves given by the selection theorem through two fixed points

$$(0, \bar{\gamma}(0)) = (0, 0), \quad (\rho, \bar{\gamma}(\rho)) = (\rho, 0).$$

The x component is set 0 just for simplifying notations. We rename the characteristic curves as

$$\gamma_0(t) := \gamma_{\tilde{\varepsilon}, 0, \bar{\gamma}(0)}(t), \quad \zeta_0(t) := \frac{d}{dt} u(i_{\gamma_0}(t)), \\ \gamma_\rho(t) := \gamma_{\tilde{\varepsilon}, \rho, \bar{\gamma}(\rho)}(t - \rho), \quad \zeta_\rho(t) := \frac{d}{dt} u(i_{\gamma_\rho}(t)).$$

Notice that $\gamma_0(t), \gamma_\rho(t)$ are tangent to $\bar{\gamma}$ respectively at times 0, ρ because they are characteristics. By (3.9), at respectively $t = 0, t = \rho$, one finds for $t \in [0, \rho]$

$$\frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \zeta_0 \geq \hat{\mathbf{g}}(i_{\bar{\gamma}}(t)) - \varepsilon \geq \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \bar{\zeta} + 2\varepsilon, \\ \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \zeta_\rho \geq \hat{\mathbf{g}}(i_{\bar{\gamma}}(t)) - \varepsilon \geq \frac{1}{\pm\sigma} \int_t^{t \pm \sigma} \bar{\zeta} + 2\varepsilon,$$

which means that the derivative $\bar{\zeta}$ of u along $\bar{\gamma}$ is lower than the ones ζ_0, ζ_ρ along γ_0, γ_ρ :

$$u(i_{\gamma_0}(t)) - u(0, 0) = \int_0^t \zeta_0 \stackrel{(3.9)}{\geq} \int_0^t \bar{\zeta} = u(i_{\bar{\gamma}}(t)) - u(0, 0), \\ u(\rho, 0) - u(i_{\gamma_\rho}(t)) = \int_t^\rho \zeta_\rho \stackrel{(3.9)}{\geq} \int_t^\rho \bar{\zeta} = u(\rho, 0) - u(i_{\bar{\gamma}}(t)).$$

This means that for t in $(0, \rho)$ one has

$$u(t, \gamma_0(t)) \geq u(t, \bar{\gamma}(t)), \quad u(t, \bar{\gamma}(t)) \geq u(t, \gamma_\rho(t)).$$

Being $f'(u)$ nonincreasing in turn

$$f'(u(t, \gamma_\rho(t))) \leq f'(u(t, \bar{\gamma}(t))) \leq f'(u(t, \gamma_0(t))).$$

Being characteristics, the functions above are just the slopes of the curves $\gamma_\rho, \bar{\gamma}, \gamma_0$:

- integrating $\bar{\gamma}, \gamma_0$ between 0 (where they coincide) and t ;
- integrating $\gamma_\rho, \bar{\gamma}$ between ρ (where they coincide) and t ;

one obtains

$$\bar{\gamma}(t) \leq \gamma_0(t), \quad \bar{\gamma}(t) \leq \gamma_\rho(t).$$

As a consequence of this and of the finite speed of propagation, γ_0 and γ_ρ must intersect in the time interval $[0, \rho]$, say at time ρ' . We can compute the value of u at

$$(\rho', \gamma_0(\rho')) = i_{\gamma_0}(\rho') = i_{\gamma_\rho}(\rho') = (\rho', \gamma_\rho(\rho'))$$

by the differential relation both on γ_0 , starting from 0, and on γ_ρ , starting from ρ : we have then

$$u(0, \gamma_0(0)) + \int_0^{\rho'} \zeta_0 = u(\rho', \gamma_0(\rho')) = u(\rho', \gamma_\rho(\rho')) = u(\rho, \gamma_\rho(\rho)) - \int_{\rho'}^\rho \zeta_\rho.$$

Comparing the the first and last term of this chain of identities one deduces

$$u(\rho, \gamma_\rho(\rho)) - u(0, \gamma_0(0)) = \int_0^{\rho'} \zeta_0 + \int_{\rho'}^\rho \zeta_\rho.$$

However, the times 0, ρ belong by construction to the set (3.9), and therefore

$$\int_0^{\rho'} \zeta_0 + \int_{\rho'}^\rho \zeta_\rho > \rho' \hat{\mathbf{g}}(i_{\bar{\gamma}}(0)) + (\rho - \rho') \hat{\mathbf{g}}(i_{\bar{\gamma}}(\rho)) - 2\varepsilon > \int_0^\rho \bar{\zeta} + \varepsilon.$$

Since the right-hand side is just $u(\rho, \gamma_\rho(\rho)) - u(0, \gamma_0(0)) + \varepsilon$, we reach a contradiction.

Substep 3.2: Inflection points. The previous step proves the statement in the connected components of $u^{-1}(\mathbb{R} \setminus N)$, where $N = \text{clos}(\text{Inf}(f))$. The assumption

$\mathcal{L}^1(N) = 0$ allows to show that inflection points do not matter. Indeed, by Theorem 30 the composition $U = u \circ i_\gamma$ is G -Lipschitz continuous: Lemma 41 below assures therefore that $u(i_\gamma)$ is differentiable with 0 derivative \mathcal{L}^1 -a.e. on $(u \circ i_\gamma)^{-1}(N)$. For every $r < s$, by the previous substep

$$\begin{aligned} u(i_\gamma(s)) - u(i_\gamma(r)) &= \int_r^s \zeta(t) dt \\ &= \int_{[r,s] \cap (u \circ i_\gamma)^{-1}(N)} \zeta(t) dt + \int_{[r,s] \setminus (u \circ i_\gamma)^{-1}(N)} \zeta(t) dt \\ &= \int_{[r,s] \cap (u \circ i_\gamma)^{-1}(N)} 0 dt + \int_{[r,s] \setminus (u \circ i_\gamma)^{-1}(N)} \hat{g}(i_\gamma(t)) dt. \end{aligned}$$

(Remember that $\hat{g} = 0$ on $u^{-1}(N)$ by definition.) This yields the thesis of Theorem 37:

$$u(i_\gamma(s)) - u(i_\gamma(r)) = \int_r^s \hat{g}(i_\gamma(t)) dt \quad \forall 0 \leq r \leq s.$$

Lemma 41. *Consider a Lipschitz continuous function $U : \mathbb{R} \rightarrow \mathbb{R}$ and a Lebesgue negligible set $N \subset \mathbb{R}$. Then the derivative of U vanishes \mathcal{L}^1 -a.e. on $U^{-1}(N)$.*

Proof. Let T_N be the set of Lebesgue points of $U^{-1}(N)$:

$$T_N = \left\{ \bar{t} : \lim_{h \downarrow 0} \frac{\mathcal{L}^1(\bar{t} - h, \bar{t} + h) \times \{\bar{y}\} \setminus U^{-1}(N))}{h} = 0 \right\}.$$

If U is G -Lipschitz continuous, then for $\bar{t} \in T_N$ one has

$$\begin{aligned} \frac{|U(\bar{t} + h) - U(\bar{t})|}{h} &\leq \frac{\mathcal{L}^1(U([\bar{t}, \bar{t} + h]) \cap N)}{h} + \frac{\mathcal{L}^1(U([\bar{t}, \bar{t} + h]) \setminus N)}{h} \\ &\leq 0 + \frac{\mathcal{L}^1(U([\bar{t}, \bar{t} + h]) \setminus N)}{h} \\ &\leq G \frac{\mathcal{L}^1([\bar{t}, \bar{t} + h] \times \{\bar{y}\} \setminus U^{-1}(N))}{h}. \end{aligned}$$

The last term converges to 0 as $h \rightarrow 0$ because $t \in T_N$. This shows that $u \circ i_\gamma(t)$ is differentiable at every $\bar{t} \in T_N$ with 0 derivative: this concludes the proof of the lemma because \mathcal{L}^1 -a.e. point of any Lebesgue measurable subset of \mathbb{R} is a Lebesgue point of the set. \square

4. DISTRIBUTIONAL SOLUTIONS ARE LAGRANGIAN SOLUTIONS

Consider a continuous distributional solution of

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g_E(t, x) \quad (1.1)$$

with $f \in C^2(\mathbb{R})$, $|g_E(t, x)| \leq G$. When inflection points of f are negligible, u is Lipschitz continuous along any characteristic curve γ of u (Theorem 30). If not, then we have cases when u is not Lipschitz continuous along some characteristics [2, §4.3]), and the points where u may not be differentiable along any characteristic curve might have positive \mathcal{L}^2 -measure.

Here we work without the assumption on inflection points. We show first that continuous distributional solutions do not dissipate entropy (Lemma 42). By approximation of the entropy, this reduces to the case of negligible inflection points, where the solution is broad and therefore Lagrangian, and it exploits the consequent BV -approximation of §2.

We show then in Lemma 45 that, given an entropy continuous distributional solution u , one can find through each point a characteristic curve $\bar{\gamma}(t)$ along which u is Lipschitz continuous. As a consequence, by §A.1 one can construct a Lagrangian parameterization and deduce that u is a Lagrangian solution.

Lemma 42. *Continuous distributional solutions of (1.1) do not dissipate entropy.*

Proof. If the closure N of the inflection points of f is negligible, then by Theorem 37 a continuous distributional solution u is a broad solution, and by Lemma 16 it is in particular a Lagrangian solution. By Corollary 28, derived from the monotone approximations of Lagrangian continuous solutions, one has then that u satisfies the entropy equality.

If the inflection points of f are not negligible, one can derive the thesis by an approximation procedure. First notice that for every entropy-entropy flux pair (which means for every function $\eta \in C^{1,1}(\mathbb{R})$ and every $q \in C^{1,1}(\mathbb{R})$ satisfying $q'(z) = \eta'(z)f'(z)$) one has the entropy equality

$$\partial_t \eta(u) + \partial_x(q(u)) = \eta'(u) g_E \quad \text{in } \mathcal{D}(\mathbb{R}^+ \times \mathbb{R} \setminus u^{-1}(N))$$

by the previous step; indeed, in the open set $\mathbb{R}^+ \times \mathbb{R} \setminus u^{-1}(N)$, where we are claiming that the PDE holds in the sense of distribution, u is valued where f does not have inflection points and therefore one can apply Corollary 28.

Consider finally a decreasing family of open sets $O_k = \cup_j (a_j^k, b_j^k) \subset \mathbb{R}$ such that

- $N = \text{clos}(\text{Infl}(f)) \subset O_k$ for $k \in \mathbb{N}$;
- $|O_k \setminus N| < 1/k$.

One can approximate η in $C^1(\mathbb{R})$ with entropies $\eta_k \in C^{1,1}(\mathbb{R})$ which are linear in O_k , for $k \in \mathbb{N}$. For every interval $(a_j^k, b_j^k) \subset O_k$ where $\eta_k(u) = c_k u$ for some $c_k \in \mathbb{R}$, for all $k \in \mathbb{N}$ one has

$$\begin{aligned} \partial_t \eta_k(u) + \partial_x(q_k(u)) &= c_k [\partial_t u + \partial_x f(u)] \\ &= c_k g_E = \eta_k'(u) g_E && \text{in } \mathcal{D}(u^{-1}((a_j^k, b_j^k))), \\ \partial_t \eta_k(u) + \partial_x(q_k(u)) &= \eta_k'(u) g_E && \text{in } \mathcal{D}(\mathbb{R}^+ \times \mathbb{R} \setminus u^{-1}(N)). \end{aligned}$$

This shows that the entropy equality holds for the entropies $\{\eta_k\}_{k \in \mathbb{N}}$. When $\eta_k(u)$, $q_k(u)$, $\eta_k'(u)$ converge uniformly to $\eta(u)$, $q(u)$, $\eta'(u)$, then the entropy equality holds also for η . \square

Corollary 43. *If u is a continuous distributional solution of (1.1), then $f'(u)$ is a continuous distributional solution of Burgers's balance law with source term $f''(u) g_E$.*

While Lemma 42 above relies on the previous results of this paper, Lemma 45 below is instead self-contained. It is however based on maximum principle, that we recall now.

Lemma 44. *Suppose u, v are entropy solutions of the PDE*

$$\begin{aligned}\partial_t u(t, x) + \partial_x(f(u(t, x))) &= g_{E_u}(t, x), \\ \partial_t v(t, x) + \partial_x(f(v(t, x))) &= g_{E_v}(t, x),\end{aligned}$$

and that for some $G \in \mathbb{R}$ there holds

$$u(t = 0, x) \leq v(t = 0, x), \quad -G \leq g_{E_u}(t, x) \leq g_{E_v}(t, x) \leq G.$$

Then $u(t, x) \leq v(t, x)$.

Proof. See the proof of Theorem 3, Page 229, [13]. Alternative approaches are the vanishing viscosity or the operator splitting, still exploiting the uniqueness of the entropy solution. \square

Lemma 45. *Suppose u is a continuous entropy solution of the PDE*

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g_E(t, x)$$

with $f \in C^2(\mathbb{R})$, $|g_E(t, x)| \leq G$. Then at each point (t, x) there exists a characteristic along which u is Lipschitz continuous.

Proof. The proof is obtained by constructing a piecewise affine approximation of the desired characteristic curve. On the two consecutive edges of the linearized curve, the Lipschitz regularity holds by the maximum principle, being an entropy solution.

Step 1: Notation. We simplify the notation changing coordinates so that we are looking for a characteristics curve throughout the point $(0, 0)$, and defined between the times $t = -1$, $t = 1$.

As the construction is local, we directly fix a square

$$Q = [-1, 1]^2.$$

Let $G = \|g_E\|_{L^\infty(Q)}$. Set $L = \|f'(u)\|_{L^\infty(Q)}$ and $M = \|f''(u)\|_{L^\infty(Q)}$. Notice that L is an upper bound for the characteristic speed $\lambda = f'(u)$ in Q . Assume e.g. $L < 1$.

Step 2: Modulus of continuity for u and λ . As u is continuous, in the compact region Q it is uniformly continuous. Let $\omega(\delta)$ denote the following modulus of continuity of u in Q :

$$\max_Q \left\{ |t' - t|, \frac{|x' - x|}{L} \right\} \leq \delta \quad \Rightarrow \quad |u(t, x) - u(t', x')| \leq \omega(\delta).$$

An analogous modulus of continuity for $\lambda = f'(u)$ is clearly given by $M\omega(\delta)$:

$$\begin{aligned}|\lambda(t, x) - \lambda(t', x')| &= |f'(u(t, x)) - f'(u(t', x'))| \\ &\leq M |u(t, x) - u(t', x')| \leq M\omega(\delta).\end{aligned}$$

Step 3: Dependency regions. Let $\delta > 0$ and $(\bar{t}, \bar{x}) \in [-1 + \delta, 1 - \delta]^2$. One draws a backward triangle of dependency for an interval of time δ , delimited from below by the segment $\bar{t} - \delta$:

$$T(\bar{t}, \bar{x}) = \{(t, x) : \bar{t} - \delta \leq t \leq \bar{t}, \bar{x} - L(\bar{t} - t) \leq x \leq \bar{x} + L(\bar{t} - t)\}.$$

Noticing that the speed of propagation λ in the rectangle

$$\left\{ (t, x) : \max \left\{ |\bar{t} - t|, \frac{|\bar{x} - x|}{L} \right\} \leq \delta \right\}$$

is, by definition of the modulus of continuity ω , bounded by $\lambda(\bar{t}, \bar{x}) \pm M\omega(\delta)$, a smaller backward triangle of dependency is given by

$$\begin{aligned}T_\delta(\bar{t}, \bar{x}) &= \{(t, x) : \bar{t} - \delta \leq t \leq \bar{t}, \\ &\quad [\lambda(\bar{t}, \bar{x}) - M\omega(\delta)](\bar{t} - t) \leq x - \bar{x} \leq [\lambda(\bar{t}, \bar{x}) + M\omega(\delta)](\bar{t} - t)\}.\end{aligned}$$

The basis of T_δ has length $2M\delta\omega(\delta)$, which is superlinear in δ .

Step 4: Comparison of u on adjacent nodes. Let $\varepsilon > 0$. The linear functions

$$u^+(t, x) = u(\bar{t}, \bar{x}) + G(t - \bar{t}) - \varepsilon, \quad u_-(t, x) = u(\bar{t}, \bar{x}) - G(t - \bar{t}) + \varepsilon$$

satisfy both

$$u^+(\bar{t}, \bar{x}) < u(\bar{t}, \bar{x}) < u_-(\bar{t}, \bar{x})$$

and the equations

$$\partial_t u_-(t, x) + \partial_x(f(u_-(t, x))) = -G, \quad \partial_t u^+(t, x) + \partial_x(f(u^+(t, x))) = G.$$

If we had either $u_-(t, x) < u(t, x)$ or $u^+(t, x) > u(t, x)$ for all x belonging to a ε -neighborhood of the basis of the small backward triangle of dependency T_δ , we would contradict the maximum principle in Lemma 44. Therefore there exists a point x_ε belonging to the interval

$$\left(\bar{x} - [\lambda(\bar{t}, \bar{x}) + M\omega(\delta)]\delta - \varepsilon, \bar{x} + [\lambda(\bar{t}, \bar{x}) + M\omega(\delta)]\delta + \varepsilon \right),$$

which is a ε -neighborhood of the basis of T_δ , where $u(t, x_\varepsilon)$ is between u^+ and u_- :

$$\begin{aligned}u(\bar{t}, \bar{x}) + G\delta + \varepsilon &= u_-(\bar{t} - \delta, x_\varepsilon) \leq u(\bar{t} - \delta, x_\varepsilon) \\ &\leq u^+(\bar{t} - \delta, x_\varepsilon) = u(\bar{t}, \bar{x}) - G\delta - \varepsilon.\end{aligned}$$

As $\varepsilon \downarrow 0$, a subsequence of $\{x_\varepsilon\}_{\varepsilon \downarrow 0}$ must converge to a point \hat{x} belonging to the basis of T_δ :

$$\begin{aligned}\text{there exists } \hat{x} &\in [\bar{x} + (\lambda(\bar{t}, \bar{x}) - M\omega(\delta))\delta, \bar{x} + (\lambda(\bar{t}, \bar{x}) - M\omega(\delta))\delta] \\ \text{such that } |u(\bar{t} - \delta, \hat{x}) - u(\bar{t}, \bar{x})| &\leq G\delta.\end{aligned}\tag{4.1}$$

Step 5: Piecewise approximation. We construct here a piecewise affine approximation of a backward characteristic through $(0, 0)$, specifying the nodes: for $k \in \mathbb{N}$ we set $(t_k, x_k) = (0, 0)$ and let

$$(t_i, x_i) := \left(-\frac{k-i}{k}, x_i \right), \quad i = 0, \dots, k-1,$$

be a point on the basis of $T(t_{i+1}, x_{i+1})$ which satisfies (4.1) where $(\bar{t}, \bar{x}) = (t_{i+1}, x_{i+1})$. Thus

$$|u(t_i, x_i) - u(t_{i-1}, x_{i-1})| \leq G(t_i - t_{i-1}) \quad \text{for } i = 0, \dots, k-1,\tag{4.2}$$

By the choice of x_{i-1} , for every k the slope

$$\lambda_{i,k} = k(x_i - x_{i-1})$$

of each segment joining (t_{i-1}, x_{i-1}) , (t_i, x_i) satisfies

$$\lambda(t_i, x_i) - M\omega(k^{-1}) \leq \lambda_{i,k} \leq \lambda(t_i, x_i) + M\omega(k^{-1}), \quad (4.3)$$

It is in particular uniformly bounded by $L + M\omega(1)$. By the Arzelà–Ascoli theorem the piecewise affine curve γ_k with edges $\{(t_i, x_i)\}_{i=0}^k$ converge uniformly as $k \uparrow \infty$, up to subsequence, to a continuous curve γ . As λ is continuous, Equation (4.3) implies that the limit curve γ is also Lipschitz continuous with slope

$$\dot{\gamma}(t) = \lambda(t, \gamma(t)).$$

This just means that we approximated a backward characteristic curve.

Step 6: Lipschitz continuity of u along the curve. For each $k \in \mathbb{N}$ set

$$u_k(t_i) := u(t_i, x_i), \quad i = 0, \dots, k,$$

and define a function $u_k(t)$ linear in each interval (t_{i-1}, t_i) , $i = 1, \dots, k$. Equation (4.2) implies that $u_k(t)$ is G -Lipschitz continuous. By the continuity of u and by the uniform convergence to γ of the piecewise affine paths γ_k with edges $\{(t_i, x_i)\}_{i=0}^k$, the function $u_k(t)$ converges uniformly to $u(i_\gamma(t))$: one has therefore that $u(i_\gamma(t))$ is itself G -Lipschitz continuous.

Step 7: Forward characteristic. We give two explanations for this step. First, Lemma 42 ensures that there is no entropy dissipation: one can thus reverse the time. Applying the above procedure for the reversed time one finds a forward characteristic curve. If one does not want to apply that strong lemma, it is enough being able to construct through each point $(t, x) \in (0, 1) \times (-1, 1)$ a backward characteristic $\gamma_{(t,x)}(s)$ along which u is G -Lipschitz continuous. Having that, the function

$$\gamma(t) := \inf \{x : \gamma_{(t,x)}(0) \geq 0\} \quad \text{with } t \in (0, 1)$$

can be verified to be a characteristic curve passing through the origin. Moreover, it is the uniform limit of characteristics along which u is G -Lipschitz continuous; in particular, by the continuity, u is therefore G -Lipschitz continuous along γ itself. \square

Corollary 46. *Suppose u is a continuous distributional solution of the PDE*

$$\partial_t u(t, x) + \partial_x(f(u(t, x))) = g_E(t, x)$$

with $\|g_E\|_{L^\infty} \leq G$. Then u is also a Lagrangian solution, with a Lagrangian source g_L bounded by G .

Proof. Lemma 42 yields that u is entropic. One can then apply Lemma 45, providing through any point a characteristic along which u is G -Lipschitz continuous. Lemmas 17, 18 finally show that u is a Lagrangian solution. \square

APPENDIX A. THREE SUFFICIENT CONDITIONS FOR THE LAGRANGIAN FORMULATION

Given a continuous function u , we consider here some sufficient conditions for satisfying the Lagrangian formulation. The section extends constructions in [5, §Appendix].

A.1. A dense set of characteristics

Fix a continuous function u . For having a Lagrangian parameterization along which u is G -Lipschitz continuous, one clearly needs through each point of the domain a characteristic curve along which u is G -Lipschitz continuous. We prove here that this is sufficient. This is Lemma 17 in the introduction.

Proof of Lemma 17. Simplify the domain to $u \in C_c(\mathbb{R}^+ \times \mathbb{R})$, as it is a local argument, and let $G > 0$. We assume that there exists a curve $\gamma_{(t,x)}(s)$ through each point (t, x) of a dense subset of $\mathbb{R}^+ \times \mathbb{R}$ such that $s \mapsto u(i_{\gamma_{(t,x)}}(s))$ is G -Lipschitz continuous. We are going to modify these characteristics in order to provide a Lagrangian parameterization.

Consider an enumeration $\{(x_{r_k}, y_{r_k})\}_{k \in \mathbb{N}}$ of a countable set of points, dense in the upper plane $\mathbb{R}^+ \times \mathbb{R}$, where the characteristic curves are given by hypothesis. We associate recursively to each point of this set a characteristic curve $\gamma_k(s)$ and we define a linear order among those:

- let $\gamma_1(s) = \gamma_{(t_{r_1}, x_{r_1})}(s)$ and write, for $k \in \mathbb{N}$,

$$r_k \preceq r_1 \text{ if } x_{r_k} \leq \gamma_1(t_{r_k}), \quad r_1 \preceq r_k \text{ if } x_{r_k} \geq \gamma_1(t_{r_k});$$

- let $h \in \mathbb{N}$ and define the new characteristic curve $\gamma_{h+1}(s)$ through $(t_{r_{h+1}}, x_{r_{h+1}})$ in order to preserve the order relation that we are establishing:

$$\gamma_{h+1}(s) = \min_{r_h \preceq r_k, k \leq h} \left\{ \gamma_k(s), \max_{r_\ell \preceq r_h, \ell \leq h} \left\{ \gamma_\ell(s), \gamma_{(t_{r_{h+1}}, x_{r_{h+1}})}(s) \right\} \right\};$$

as $\gamma_k(s)$, for $k \leq h$, and $\gamma_{(t_{r_{h+1}}, x_{r_{h+1}})}(s)$ are characteristic curves along which u is G -Lipschitz continuous by hypothesis, then also $\gamma_{h+1}(s)$ is a characteristic curve along which u is G -Lipschitz continuous;

- we set then, for $k \in \mathbb{N}$,

$$r_k \preceq r_{h+1} \text{ if } x_{r_k} \leq \gamma_{h+1}(t_{r_k}), \quad r_{h+1} \preceq r_k \text{ if } x_{r_k} \geq \gamma_{h+1}(t_{r_k});$$

by construction this extends the relation defined in the previous items.

The set of uniformly Lipschitz continuous curves

$$\mathcal{C} = \{\gamma_k(s)\}_{k \in \mathbb{N}}$$

is totally ordered and the images of these curves are dense in $\mathbb{R}^+ \times \mathbb{R}$. We can complete this set in the uniform topology: the curves that we introduce with the closure are still characteristic curves because of the continuity of $f'(u)$; as well, u is G -Lipschitz continuous along them and they preserve the order, in the sense that

any two curves do not cross each other but always lie on a fixed side, when they differ. If $\{q_k\}_{k \in \mathbb{N}}$ is an enumeration of the rational numbers, the map

$$\begin{aligned} \theta : \text{clos}(\mathcal{C}) &\rightarrow \mathbb{R} \\ \gamma &\mapsto \sum_{k=0}^{\infty} \frac{\gamma(q_k)}{2^{-k}} \end{aligned}$$

is continuous and strictly order preserving. In particular, it is invertible with continuous inverse.

One can then verify that a Lagrangian parameterization is provided by

$$\chi(s, y) = [\theta^{-1}(y)](s) \quad \text{for } s \in \theta(\text{clos}(\mathcal{C})).$$

By construction $t \mapsto U(t, y) = u(t, \chi(t, y))$ is G -Lipschitz continuous for each y fixed: the thesis thus follows by Lemma 18. \square

A.2. Lipschitz continuity along characteristics

Fix a continuous function u . For having that u is a Lagrangian solution, one clearly needs that $t \mapsto u(t, \chi(t, y))$ is Lipschitz continuous, uniformly in the y parameter, for some Lagrangian parameterization χ . We prove here that this is sufficient. This is Lemma 18 in the introduction.

We are not concerned here with the compatibility of the source terms.

Proof of Lemma 18. Simplify the domain to $u \in C_c(\mathbb{R}^+ \times \mathbb{R})$, as it is a local argument, and let $G > 0$. We want to show that if there exists a Lagrangian parameterization χ such that

$$-G \leq \frac{d}{dt} u(t, \chi(t, y)) \leq G \quad \text{in } \mathcal{D}(i_{\chi(y)}^{-1}(\Omega)) \text{ for all } y \in \mathbb{R},$$

then one can find a function $g_L \in \mathfrak{L}_L^\infty(\Omega)$ such that

$$\frac{d}{dt} u(t, \chi(t, y)) = g_L(t, \chi(t, y)) \quad \text{in } \mathcal{D}(i_{\chi(y)}^{-1}(\Omega)) \text{ for all } y \in \mathbb{R}.$$

Set $U(t, y) = u(t, \chi(t, y))$ and consider $\mathfrak{G} \in \mathfrak{L}^\infty(\mathbb{R}^+ \times \mathbb{R})$ such that, in the (t, y) -half plane,

$$\mathfrak{G}(t, y) = \partial_t U(t, y) \quad \text{in } \mathcal{D}(i_{\chi(y)}^{-1}(\Omega)) \text{ for all } y \in \mathbb{R}.$$

We want to show that it can be chosen of the form $\mathfrak{G}(t, y) = g_L(t, \chi(t, y))$ for some g_L , which means that it is essentially single-valued on the level sets of χ . Fixed y , we show the following: the set of times t where u has a Lebesgue point of classical differentiability

- (i) both along the characteristic curve $\gamma(t) = \chi(t, y)$,
- (ii) and also along another characteristic curve $\tilde{\gamma}(t)$, lying on a fixed side of $\gamma(t)$,
- (iii) with two different values of the derivative,

is at most countable. This is enough since characteristics of a same Lagrangian parameterization are by definition ordered.

Let $\varepsilon, \sigma > 0$. By a reduction argument it suffices to show the following *claim*: let $S(y)$ be the set of all t such that

$$\begin{aligned} &\left| \frac{U(t+h, y) - U(t, y)}{h} - \mathfrak{G}(t, y) \right| < \varepsilon, \\ &\text{there exists } \gamma(s) \text{ characteristic with } \gamma(t) = \chi(t, y), \\ &\text{and for every } |h| \leq \sigma \text{ there holds } \gamma(t+h) \leq \chi(t+h, y), \\ &\frac{u(t+h, \gamma(t+h)) - u(t, \gamma(t))}{h} - \mathfrak{G}(t, y) > \varepsilon, \end{aligned} \tag{A.1}$$

then $S(y)$ does not contain two points t_1, t_2 closer than σ . Indeed, if we are comparing the value of the derivative of u along different characteristics of the parameterization χ , then an order condition is satisfied among characteristics. Moreover, as we consider Lebesgue points of differentiability, with different values for the derivative of u along $\chi_y^{\text{el}}(s)$ and $\gamma(s)$, up to a countable covering we are dealing with sets like (A.1).

We prove the claim by contradiction: let

$$t_1, t_2 \in S(y) \text{ with } 0 < t_2 - t_1 < \sigma.$$

The definition (A.1) of $S(y)$ provides curves γ_1, γ_2 which intersect

$$\gamma_0(s) := \chi(s, y)$$

at times t_1, t_2 respectively, and which for $t_1 \leq s \leq t_2$ satisfy the additional properties

$$\begin{aligned} \gamma_1(s) &\leq \gamma_0(s), & u(s, \gamma_1(s)) - u(t_1, \gamma_1(t_1)) &> (\mathfrak{G}(t, y) + \varepsilon)(s - t_1), \\ \gamma_2(s) &\leq \gamma_0(s), & u(t_2, \gamma_2(t_2)) - u(s, \gamma_2(s)) &> (\mathfrak{G}(t, y) + \varepsilon)(t_2 - s). \end{aligned}$$

By the ordering imposed in (A.1) and by the uniform Lipschitz continuity implied by the fact that they are characteristics, the curves $\gamma_1(s), \gamma_2(s)$ necessarily meet at some time $\bar{t} \in [t_1, t_2]$. One can then compute the difference $U(t_2, y) - U(t_1, y)$ in two ways:

- by applying the incremental relation in (A.1) relative to χ , which gives

$$\begin{aligned} U(t_2, y) - U(t_1, y) &= U(t_2, y) - U(\bar{t}, y) + U(\bar{t}, y) - U(t_1, y) \\ &< \mathfrak{G}(t_2, y)(t_2 - \bar{t}) + \mathfrak{G}(t_1, y)(\bar{t} - t_1) + 2\varepsilon; \end{aligned}$$
- by applying the incremental relation in (A.1) relative to γ_1, γ_2 : denoting by \bar{x} the value $\gamma_1(\bar{t}) = \gamma_2(\bar{t})$ when γ_1 and γ_2 intersect one has

$$\begin{aligned} U(t_2, y) - U(t_1, y) &= u(t_2, \gamma_2(t_2)) - u(\bar{t}, \bar{x}) + u(\bar{t}, \bar{x}) - u(t_1, \gamma_1(t_1)) \\ &> \mathfrak{G}(t_2, y)(t_2 - \bar{t}) + \mathfrak{G}(t_1, y)(\bar{t} - t_1) + 2\varepsilon. \end{aligned}$$

The estimates that we obtain in these two ways are not compatible: we have reached a contradiction. \square

A.3. Stability of the Lagrangian formulation for uniform convergence of u

We state for completeness that Lagrangian solutions are closed w.r.t. uniform convergence, provided the sources are uniformly bounded. We include this for completeness but it follows easily by the previous analysis of the section.

Corollary 47. Let $G > 0$ and $u_k(t, x)$ be a sequence of continuous Lagrangian solutions of

$$\partial_t u_k(t, x) + \partial_x(f(u_k(t, x))) = g_{E_k}(t, x) \quad (1.1)$$

with $f \in C^2(\mathbb{R})$, $|g_{E_k}(t, x)| \leq G$. If u_k converges uniformly to u , then u is a Lagrangian solution with source term bounded by G .

Proof. We verify that through every point $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$ there exists a characteristic curve $\gamma(s)$ such that $u(i_\gamma(s))$ is G -Lipschitz continuous: Lemma 17 then provides a Lagrangian parameterization along which u is G -Lipschitz continuous, and Lemma 18 gives the thesis.

As $\{u_k\}_{k \in \mathbb{N}}$ are Lagrangian solutions of (1.1) with sources uniformly bounded by G , one can find for each $k \in \mathbb{N}$ a characteristic curve $\gamma_k(s)$ of u_k through (t, x) satisfying

$$|u_k(i_{\gamma_k}(r)) - u_k(i_{\gamma_k}(s))| \leq G|r - s|. \quad (A.2)$$

The family $\{\gamma_k(s)\}_{k \in \mathbb{N}}$ is locally equi-Lipschitz continuous and equi-bounded, as $\gamma_k(t) = x$. By the Arzelà–Ascoli theorem this family has a subfamily uniformly convergent to a function $\gamma(s)$. From the uniform convergence of the continuous functions u_k and γ_k , the relation

$$\gamma_k(r) - \gamma_k(s) = \int_r^s u_k(i_{\gamma_k}(q)) dq$$

goes to the limit and it implies that γ is characteristic curve for u . Moreover, also (A.2) goes to the limit and it yields that u is G -Lipschitz continuous along γ . \square

NOMENCLATURE

- $[\cdot]_\lambda, [\cdot]_\chi, [\cdot]$ projections on $\mathfrak{L}_B^\infty(X)$, $\mathfrak{L}_L^\infty(X)$, $L^\infty(X)$ respectively, Notation 9,
- χ Lagrangian parameterization for a continuous solution u to (1.1), Definition 5,
- $\text{clos}(\cdot)$ closure of a set,
- $\mathcal{D}(\Omega)$ distributions on Ω , Notation 8,
- $L^\infty(X)$ bounded functions on X , identified \mathcal{L}^2 -a.e., Notation 8,
- D_t, D_x partial derivatives of a function of bounded variation, Notation 3,
- γ, i_γ characteristic curve, Definition 4,
- g, g_B, g_L functions belonging to $\mathfrak{L}^\infty(X)$, $\mathfrak{L}_B^\infty(X)$, $\mathfrak{L}_L^\infty(X)$ respectively, Notation 8,
- g, g_E distributional, bounded source term for the balance law (1.1),
- $\text{Infl}(f)$ inflection points of f , Definition 15,
- λ the composite function $f' \circ u$, Notation 1,
- $\mathcal{L}^1, \mathcal{L}^2$ 1- and 2-dimensional Lebesgue measure,
- $\mathcal{M}(X)$ Radon measures on X , Notation 8,
- Ω open subset of $\mathbb{R}^+ \times \mathbb{R}$ (connected if needed),

- $\frac{\partial}{\partial t}, \frac{\partial}{\partial x}$ classical partial derivatives, Notation 3,
- ∂_t, ∂_x distributional partial derivatives, Notation 3,
- $\mathfrak{L}^\infty(X)$ functions defined pointwise on X , Notation 8,
- $\mathfrak{L}_B^\infty(X)$ functions coinciding \mathcal{L}^1 -a.e. on characteristics of u , Notation 8,
- $\mathfrak{L}_L^\infty(X)$ functions coinciding \mathcal{L}^1 -a.e. on the Lagrangian parameterization χ , Notation 8,
- $\varphi_t^{e_2}(x)$ restriction of a function $\varphi(t, x)$ to the second coordinate, Notation 2,
- $\varphi_x^{e_1}(t)$ restriction of a function $\varphi(t, x)$ to the first coordinate, Notation 2,
- $C(\Omega)$ continuous functions on Ω , see also $C_b, C^k, C_c^k, C^{k,1/\alpha}$ in Notation 8,
- f flux function for the balance law (1.1),
- u continuous solution, Notation 1,
- X subset of $\mathbb{R}^+ \times \mathbb{R}$, usually Borel.

ACKNOWLEDGEMENTS

The authors wish to thank Francesco Bigolin and Francesco Serra Cassano, from the University of Trento, for interesting discussions which further motivated this later paper. We are grateful to the anonymous referee for interesting questions improving our bibliographical references. This work was supported by the Centro di Ricerca Matematica Ennio De Giorgi (Pisa), the EPSRC Science and Innovation award to OxPDE (EP/E035027/1), the Istituto Nazionale di Alta Matematica (INdAM), and the PRIN grant “Nonlinear hyperbolic partial differential equations, dispersive and transport equations: theoretical and applicative aspects”.

REFERENCES

- [1] ALBERTI, GIOVANNI; BIANCHINI, STEFANO; CARAVENNA, LAURA. Reduction on characteristics for continuous solution of a scalar balance law. *Hyperbolic problems: theory, numerics, applications*. Proceedings of the 14th International Conference on Hyperbolic Problems (HYP2012), Padua, June 25-29, 2012. Edited by F. Ancona et al., pp. 399–406. AIMS Series on Applied Mathematics, vol. 8. American Institute of Mathematical Sciences, Springfield (MO), 2014.
- [2] ALBERTI, GIOVANNI; BIANCHINI, STEFANO; CARAVENNA, LAURA. Eulerian, Lagrangian and Broad continuous solutions to a balance law with non-convex flux. II. Preprint SISSA 32/2016/MATE.
- [3] AMBROSIO, LUIGI; FUSCO, NICOLA; PALLARA, DIEGO. Functions of bounded variation and free discontinuity problems. Oxford Mathematical Monographs. Clarendon Press, Oxford, 2000.
- [4] BIGOLIN, FRANCESCO; SERRA CASSANO, FRANCESCO. Intrinsic regular graphs in Heisenberg groups vs. weak solutions of non-linear first-order PDEs. *Adv. Calc. Var.*, 3 (2010), no. 1, 69–97.
- [5] BIGOLIN, FRANCESCO; CARAVENNA, LAURA; SERRA CASSANO, FRANCESCO. Intrinsic Lipschitz graphs in Heisenberg groups and continuous solutions of a balance equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 32 (2015), no. 5, 925–963.
- [6] BRESSAN, ALBERTO. *Hyperbolic systems of conservation laws. The one-dimensional Cauchy problem*. Oxford Lecture Series in Mathematics and its Applications, vol. 20. Oxford University Press, Oxford, 2000.

- [7] CITTI, GIOVANNA; MANFREDINI, MARIA; PINAMONTI, ANDREA; SERRA CASSANO, FRANCESCO. Smooth approximation for intrinsic Lipschitz functions in the Heisenberg group. *Calc. Var. Partial Differential Equations*, 49 (2014), no. 3-4, 1279–1308.
- [8] DAFERMOS, CONSTANTINE M. Continuous solutions for balance laws. *Ric. Mat.*, 55 (2006), no. 1, 79–91.
- [9] DE GIORGI, ENNIO; COLOMBINI, FERRUCCIO; PICCININI, LIVIO C. *Frontiere orientate di misura minima e questioni collegate*. Pubblicazioni della Classe di Scienze. Scuola Normale Superiore, Pisa, 1972.
- [10] FEDERER, HERBERT. *Geometric measure theory*. Grundlehren der mathematischen Wissenschaften, vol. 153. Springer Verlag, Berlin-New York, 1969. Reprinted in the series Classics in Mathematics. Springer Verlag, Berlin-Heidelberg, 1996.
- [11] FRANCHI, BRUNO; SERAPIONI, RAUL; SERRA CASSANO, FRANCESCO. Differentiability of intrinsic Lipschitz functions within Heisenberg groups. *J. Geom. Anal.* 21 (2011), no. 4, 1044–1084.
- [12] KIRCHHEIM, BERND; SERRA CASSANO, FRANCESCO. Rectifiability and parameterization of intrinsic regular surfaces in the Heisenberg group. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5)*, 3 (2004), no. 4, 871–896.
- [13] KRUŽKOV, STANISLAV N. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*, 81(123) (1970), 228–255. English translation in *Math. USSR Sb.*, 10 (1970), 217–243.
- [14] KUNUGUI, KINJIRO. Contributions à la théorie des ensembles boreliens et analytiques, II and III. *J. Fac. Sci. Hokkaido Imp. Univ. Ser. I.*, 8 (1939-1940), no. 1, 1–24, and no. 2-3, 79–108.
- [15] MONTI, ROBERTO; VITTONI, DAVIDE. Sets with finite \mathbb{H}^n -perimeter and controlled normal. *Math. Z.*, 270 (2012), no. 1-2, 351–367.
- [16] VON NEUMANN, JOHN. On rings of operators. Reduction theory. *Ann. of Math. (2)*, 50 (1949), 401–485.
- [17] ROŽDESTVENSKIĪ, BORIS L.; YANENKO, NIKOLAI N. *Systems of quasilinear equations and their applications to gas dynamics*. Translations of Mathematical Monographs, vol. 55. American Mathematical Society, Providence (RI), 1983.
- [18] SRIVASTAVA, SHASHI MOHAN. *A course on Borel sets*. Graduate Texts in Mathematics, vol. 180. Springer-Verlag, New York, 1998.
- [19] VITTONI, DEVIDE. *Submanifolds in Carnot groups*. Tesi della Scuola Normale Superiore di Pisa (nuova serie) [Theses of Scuola Normale Superiore di Pisa (new series)], vol. 7. Edizioni della Normale, Pisa, 2008.

G.A.

Dipartimento di Matematica, Università di Pisa, largo Pontecorvo 5, 56127 Pisa, Italy
e-mail: giovanni.alberti@unipi.it

S.B.

SISSA-ISAS, via Bonomea 265, 34136 Trieste, Italy
e-mail: stefano.bianchini@sissa.it

L.C.

Dipartimento di Matematica, Università di Padova, via Trieste 63, 35121 Padova, Italy
e-mail: laura.caravenna@unipd.it