

REGULARITY THEOREMS, UP TO THE BOUNDARY, FOR SHEAR THICKENING FLOWS[☆]

H. Beirão da Veiga^a, P. Kaplický^b, M. Růžička^c

^a*Dipartimento di Matematica Applicata "U.Dini", Via Buonarroti, 1/C, I-56127 Pisa, Italy*

^b*Charles University in Prague, Sokolovská 83, Praha 8, 18675, Czech Republic*

^c*Mathematical Institut, University Freiburg, Eckerstr. 1, D-79104 Freiburg, Germany*

Abstract

This article concerns the regularity up to the boundary of weak solutions to systems describing the flow of generalized Newtonian shear thickening fluids under the homogeneous Dirichlet boundary condition. The extra stress tensor $\mathcal{S}(D)$, see (1.2) below, is given by a power law with shear exponent $p \geq 2$. Complete proofs of the results presented here are given in the forthcoming paper [4]. The aim of this note is to describe the results proved in [4], together with suitable comments.

Résumé

Théorèmes de régularité, jusqu'au bord, pour des fluides à viscosité dilatante.

Cet article regarde la régularité jusqu'au bord des solutions faibles de systèmes décrivant le mouvement de fluides Newtonian généralisés, à viscosité dilatante, avec des conditions de Dirichlet à la frontière. Le tenseur des tensions supplémentaires $\mathcal{S}(D)$, voir (1.2), est donné par une loi de puissance avec exposant $p \geq 2$. Démonstrations complètes des résultats présentés ici sont donnés dans l'article [4], à paraître. Le but de cette note est de décrire les résultats démontrés dans [4], avec des commentaires adéquats.

1. Introduction

This note is concerned with the system

$$\begin{aligned}\partial_t u - \operatorname{div} \mathcal{S}(Du) + [\nabla u]u + \nabla \pi &= f, \\ \operatorname{div} u &= 0,\end{aligned}\tag{1.1}$$

[☆]Michael Růžička has been supported by DFG Forschergruppe "Nonlinear Partial Differential Equations: Theoretical and Numerical Analysis". Hugo Beirão da Veiga and Petr Kaplický thank the University of Freiburg for the kind hospitality during part of the preparation of the paper. Research of Petr Kaplický was also supported by the grant GACR 201/09/0917 and partially also by the research project MSM 0021620839.

Email addresses: bveiga@dma.unipi.it (H. Beirão da Veiga), kaplicky@karlin.mff.cuni.cz (P. Kaplický), rose@mathematik.uni-freiburg.de (M. Růžička)

describing the motion of a fluid in $\Omega \times I$, where $\Omega \subset \mathbb{R}^n$ and I is interval $(0, T)$. The unknowns are the velocity $u : \Omega \times I \rightarrow \mathbb{R}^n$ and the pressure $\pi : \Omega \times I \rightarrow \mathbb{R}$. The external force f is given, and the convective term is defined as $([\nabla u]u)_i = \sum_{j=1}^n u_j \partial_j u_i$, $i = 1, \dots, n$. The extra stress tensor \mathcal{S} , expressing the inner properties of the fluid, depends only on $Du = (\nabla u + \nabla u^T)/2$ by the principle of material frame indifference, and is assumed to possess p -structure with $p \geq 2$. Under these assumptions the system (1.1) models the flow of shear thickening fluids. Typical examples are

$$\mathcal{S}(D) = (1 + |D|^2)^{\frac{p-2}{2}} D \quad \text{or} \quad \mathcal{S}(D) = (1 + |D|)^{p-2} D. \quad (1.2)$$

We assume the non-slip boundary condition on $\partial\Omega \times I$

$$u = 0. \quad (1.3)$$

We do not impose an initial condition since the results are local in time.

The system (1.1) is nowadays classical. It was proposed by O.A. Ladyzhenskaya (see [11] and references) as a modification of the Navier–Stokes system. J.-L. Lions (see [12]), suggested a similar system, however with the elliptic term depending on the full gradient. In spite of the fact that the system has been extensively studied, there are still many open problems, especially concerning the regularity of weak solutions. Our results contribute to the field of regularity properties of weak solutions to the problems (1.1) and its steady version (1.4). In particular, we prove boundary regularity of weak solutions (u, π) of the problem (1.1), (1.2), and (1.3) for suitable $p \geq 2$. The main obstacles arise close to the boundary. They are connected with the structure of the elliptic term, namely that it is nonlinear and depends on the symmetric part of the gradient only, and with the presence of pressure in the equation. This is why we do not get the same regularity as for solutions of Stokes problem. The difficulties due to convective term are avoided by considering sufficiently large $p \geq 2$.

The question of regularity of weak solutions of (1.1) is transferred, by appealing to [6], to the problem of the regularity of weak solutions to the stationary variant of (1.1) in Ω , namely

$$\begin{aligned} -\operatorname{div} \mathcal{S}(Du) + \delta [\nabla u]u + \nabla \pi &= f, \\ \operatorname{div} u &= 0, \end{aligned} \quad (1.4)$$

where $\delta = 0$ or 1 . Since the problem with $\delta = 0$ already contains the principle features that appear due to the form of \mathcal{S} , we treat the regularity of solutions to (1.4) with $\delta = 0$ as the basic step of the proof.

Let us now formulate precisely the assumptions on \mathcal{S} .

Assumption 1 (extra stress tensor). *We assume that the extra stress tensor $\mathcal{S} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}_{\text{sym}}^{n \times n}$ belongs to $C^0(\mathbb{R}^{n \times n}, \mathbb{R}_{\text{sym}}^{n \times n}) \cap C^1(\mathbb{R}^{n \times n} \setminus \{0\}, \mathbb{R}_{\text{sym}}^{n \times n})$, where $\mathbb{R}_{\text{sym}}^{n \times n} := \{A \in \mathbb{R}^{n \times n} \mid A = A^\top\}$, and satisfies $\mathcal{S}(0) = 0$ and $\mathcal{S}(A) = \mathcal{S}(A^{\text{sym}})$, where $A^{\text{sym}} := \frac{1}{2}(A + A^\top)$. Moreover, we assume that \mathcal{S} has p -structure, i.e., there exists $p \in (1, \infty)$ such that*

$$\sum_{i,j,k,l=1}^n \partial_{kl} \mathcal{S}_{ij}(A) B_{ij} B_{kl} \geq c (1 + |A^{\text{sym}}|)^{p-2} |B^{\text{sym}}|^2, \quad (1.5a)$$

$$|\partial_{kl} \mathcal{S}_{ij}(A)| \leq C (1 + |A^{\text{sym}}|)^{p-2} \quad (1.5b)$$

is satisfied for all $A, B \in \mathbb{R}^{n \times n}$ with $A^{\text{sym}} \neq 0$.

These assumptions are motivated by the typical prototypes for the extra stress tensor given in (1.2). We refer the reader to [13], [8], [5] for a more detailed discussion leading to Assumption 1. We also define a function $\mathcal{F}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}_{\text{sym}}^{n \times n}$, closely related to the extra stress tensor \mathcal{S} , by

$$\mathcal{F}(A) := (1 + |A^{\text{sym}}|)^{\frac{p-2}{2}} A^{\text{sym}}. \quad (1.6)$$

Note that $\mathcal{F}(D) \sim |D|^{p/2}$. Since in the following and in reference [4] we insert into \mathcal{S} and \mathcal{F} only symmetric tensors, we can drop the superscripts “sym” and restrict the admitted tensors to symmetric ones.

2. Main results

Theorem 2.1. *Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded domain with $C^{2,1}$ boundary, let \mathcal{S} satisfy Assumption 1 with $p \geq 2$, and let $f \in L^2(\Omega)$. Then the unique weak solution $u \in W_0^{1,p}(\Omega)$, with $\text{div } u = 0$, of problem (1.4), with $\delta = 0$, and boundary condition (1.3) satisfies*

$$u \in W^{1,q}(\Omega), \quad \mathcal{F}(Du) \in W^{1, \frac{2q}{p+q-2}}(\Omega), \quad (2.1)$$

for $q = (np + 2 - p)/(n - 2)$ if $n \geq 3$, and for all $q < +\infty$ if $n = 2$.

Theorem 2.2. *Let Ω and f be as in Theorem 2.1, and let \mathcal{S} satisfy Assumption 1 with $p \geq \max\{2, (3n)/(n+2)\}$. Then, for $p \in [3, \infty) \cup (n/2, \infty)$, any weak solution $u \in W_0^{1,p}(\Omega)$, with $\text{div } u = 0$, of problem (1.4), with $\delta = 1$, and boundary condition (1.3) satisfies (2.1), where q is as above.*

Remark 2.1. (i) *In the interior and in tangential directions we get better regularity properties in the previous theorems. More precisely*

$$\mathcal{F}(Du) \in W_{\text{loc}}^{1,2}(\Omega), \quad \xi \partial_\tau \mathcal{F}(Du) \in L^2(\Omega),$$

where ξ is some cut off function with support near the boundary $\partial\Omega$ and the tangential derivative ∂_τ is defined in a suitable neighborhood of the boundary.

(ii) *One has*

$$|\nabla \mathcal{F}(Du)| \simeq (1 + |Du|)^{\frac{p-2}{2}} |\nabla Du|$$

and also $|\partial_\tau \mathcal{F}(Du)| \simeq (1 + |Du|)^{\frac{p-2}{2}} |D \partial_\tau u|$. For $p \geq 2$ it follows that

$$|\nabla^2 u| \leq c |\nabla \mathcal{F}(Du)|.$$

Theorem 2.3. *Let \mathcal{S} satisfy Assumption 1 with $p > (3n+2)/(n+2)$. Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded domain with $C^{2,1}$ boundary, $I = (0, T)$, $T > 0$, and $f \in W_{\text{loc}}^{1,2}(I, L^2(\Omega))$. Then any weak solution $u \in L^\infty(I, L^2(\Omega)) \cap L^p(I, W_0^{1,p}(\Omega))$, with $\text{div } u = 0$, of the problem (1.1), and boundary condition (1.3) satisfies*

$$\begin{aligned} u &\in W_{\text{loc}}^{1,\infty}(I, L^2(\Omega)), \quad \mathcal{F}(Du) \in W_{\text{loc}}^{1,2}(I, L^2(\Omega)) \\ u &\in L_{\text{loc}}^\infty(I, W^{1,q}(\Omega)), \quad \mathcal{F}(Du) \in L_{\text{loc}}^\infty(I, W^{1, \frac{2q}{p+q-2}}(\Omega)) \end{aligned} \quad (2.2)$$

for $q = (np + 2 - p)/(n - 2)$, ($q < +\infty$ if $n = 2$).

Remark 2.2. *Note that Remark 2.1 also applies to Theorem 2.3. In particular we get*

$$\mathcal{F}(Du) \in W_{\text{loc}}^{1,2}(I \times \Omega), \quad \xi \partial_\tau \mathcal{F}(Du) \in L_{\text{loc}}^\infty(I, L^2(\Omega)).$$

3. Further remarks

The main obstacle in the proofs of the above theorems is the boundary condition prescribed on the non-flat boundary $\partial\Omega$ together with the fact that the extra stress tensor \mathcal{S} depends only on symmetric part of the velocity gradient. Moreover, the incompressibility of the flow (which results in the fact that weak solutions are divergence free, and in the appearance of the pressure term $\nabla\pi$) causes additional problems. Let us now briefly describe how we treat these difficulties in the case of the steady problem with $\delta = 0$.

The regularity of the tangential derivatives $\partial_\tau u$ near the boundary, namely

$$\int_{\Omega} \xi |\partial_\tau \mathcal{F}(Du)|^2 dx \simeq \int_{\Omega} \xi (1 + |Du|)^{p-2} |D\partial_\tau u|^2 dx < C, \quad (3.1)$$

is obtained by the classical difference quotients method. We appeal to translations parallel to the non-flat boundary $\partial\Omega$. When deriving the main estimates there appear some terms which are not present if the boundary is flat. The normal derivatives are restored from the equations (1.1). Due to the smoothness of $\partial\Omega$, it is possible to express the whole second gradient of u via the gradient of the tangential derivatives of u and π . Here, in appealing to (3.1), one has to take into account that we do not know if, for $p \neq 2$, a Korn's inequality, of the type

$$\int_{\Omega} (1 + |Du|)^{p-2} |\nabla \partial_\tau u|^2 dx \leq C \int_{\Omega} (1 + |Du|)^{p-2} |D\partial_\tau u|^2 dx,$$

holds. This situation leads to a lower regularity in the normal direction.

The results in this article are new and improve all the previous results. They are obtained by combining several methods from [10], [14], [1], [3] with some new ideas. As in [10], [14], and [2], we consider non-flat boundaries. In particular, we improve the regularity exponents obtained in [2] by finding a better balance between the two main terms which prevent optimal results.

The treatment of the unsteady problem (1.1) is based on the results for the steady problem (1.4) and on the improvements of the time regularity of weak solutions shown in [6].

Apart from the results mentioned above, there are some other investigations dealing with the boundary regularity of weak solutions to systems related to (1.1). For some references we refer the reader to [4]. We just mention here the forthcoming paper [7], where the authors prove, for small data f , that the solutions belong to $C^{1,\alpha}(\bar{\Omega}) \cap W^{2,2}(\Omega)$, up to the boundary. Finally we claim that the method followed in [4] may lead to an improvement of the result in [9] for all $p \geq 2$.

References

- [1] H. BEIRÃO DA VEIGA, *On the regularity of flows with Ladyzhenskaya shear-dependent viscosity and slip or nonslip boundary conditions*, Comm. Pure Appl. Math. **58** (2005), no. 4, 552–577.
- [2] H. BEIRÃO DA VEIGA, *On the Ladyzhenskaya–Smagorinsky turbulence model of the Navier–Stokes Equations in Smooth Domains. The Regularity Problem*, J. Eur. Math. Soc., **11** (2009), 127–167.
- [3] H. BEIRÃO DA VEIGA, *Turbulence Models, p -fluid flows, and $W^{2,l}$ -regularity of solutions*, Comm. Pure Appl. Anal., **8** (2009), 769–783.
- [4] H. BEIRÃO DA VEIGA, P. KAPLICKÝ, AND M. RŮŽIČKA, *Boundary regularity of shear thickening flows*, J. Math. Fluid Mech., DOI: 10.1007/s00021-010-0025-y. Preprint series of the Charles University, Prague, MATH-kma-2008/277 340.8.

- [5] L.C. BERSELLI, L. DIENING, AND M. RŮŽIČKA, Existence of Strong Solutions for Incompressible Fluids with Shear Dependent Viscosities, *J. Math. Fluid Mech.*, in press. DOI 10.1007/s00021-008-0277-y.
- [6] M. BULÍČEK, F. ETTWEIN, P. KAPLICKÝ, AND D. PRAŽÁK , *On uniqueness and time regularity of flows of power-law like non-Newtonian fluids*, *Math. Methods Appl. Sc.*, DOI:10.1002/mma.1314 (2010)
- [7] F. CRISPO AND C. GRISANTI, *On the $C^{1,\gamma}(\overline{\Omega}) \cap W^{2,2}(\Omega)$ regularity for a class of electro-rheological fluids*, *J. Math. Anal. Appl.* **356** (2009), 119-132.
- [8] L. DIENING AND M. RŮŽIČKA, *Strong Solutions for Generalized Newtonian Fluids*, *J. Math. Fluid Mech.* **7** (2005), 413–450.
- [9] P. KAPLICKÝ, *Regularity of flows of a non-Newtonian fluid subject to Dirichlet boundary conditions*, *Z. Anal. Anwendungen* **24** (2005), no. 3, 467–486.
- [10] P. KAPLICKÝ, J. MÁLEK, AND J. STARÁ, *$C^{1,\alpha}$ -Regularity of Weak Solutions to a Class of Nonlinear Fluids in Two Dimensions - Stationary Dirichlet Problem*, *Zap. Nauchn. Sem. Pt. Odel. Mat. Inst.* **259** (1999), 89–121.
- [11] O.A. LADYZHENSKAYA, *The Mathematical Theory of Viscous Incompressible Flow*, Gordon and Breach, New York, 1969, 2nd edition.
- [12] J.L. LIONS, *Quelques Méthodes de Résolution des Problèmes aux Limites Non Linéaires*, Dunod, Paris, 1969.
- [13] J. MÁLEK, J. NEČAS, M. ROKYTA, AND M. RŮŽIČKA , *Weak and Measure-valued Solutions to Evolutionary PDEs*, *Applied Mathematics and Mathematical Computations*, vol. 13, Chapman & Hall, London, 1996.
- [14] J. MÁLEK, J. NEČAS, AND M. RŮŽIČKA , *On weak solutions to a class of non-Newtonian incompressible fluids in bounded three-dimensional domains: the case $p \geq 2$* , *Adv. Differential Equations* **6** (2001), no. 3, 257–302.