

ACCADEMIA NAZIONALE DEI LINCEI

Estratto dai Rendiconti della Classe di Scienze fisiche, matematiche e naturali

Serie VIII, vol. LVIII, fasc. 4 - Aprile 1975

Matematica. — A remark on the differentiability for Green's operators of variational inequalities. Nota di HUGO BEIRÃO DA VEIGA (*), presentata (***) dal Corrisp. G. STAMPACCHIA.

RIASSUNTO. — È stato dimostrato in [1] che l'operatore P definito da (3) è differenziabile nell'origine, inteso come operatore da L^2(Omega) in L^2(Omega). In questa Nota si osserva che continua a sussistere lo stesso risultato se P viene inteso come operatore da L^2(Omega) in W^{1,2}(Omega) ed inoltre come quest'ultimo possa essere ulteriormente generalizzato.

This Note is concerned with the recent paper [1] to which the reader is referred for terminology, notation and further details.

Let Omega be an open bounded set in the n-dimensional Euclidean space R^n and let Gamma be the boundary of Omega. We assume that Omega and Gamma are smooth.

We denote by || ||_p and || ||_{s,p} the usual norms in the space L^p(Omega) and W^{s,p}(Omega) respectively, and we put H = L^2(Omega), || || = || ||_2. We shall consider also the spaces L^p(Gamma) and W^{s,p}(Gamma) with the usual norms | | _p and | | _{s,p} respectively.

Let now alpha : R -> 2^R and suppose that 0 in alpha(0); we say that the graph alpha is differentiable at the origin, with finite derivative alpha', if the following condition holds:

- (1) for any epsilon > 0 there exists delta_epsilon > 0 such that |z - alpha'y| <= epsilon |y|, for all z in alpha(y), for all y in] - delta_epsilon, delta_epsilon [cap D(alpha).

We say that alpha is differentiable at the origin with alpha' = + infinity if

- (2) for any epsilon > 0 there exists delta_epsilon > 0 such that |y| <= epsilon |z|, for all z in alpha(y), for all y in] - delta_epsilon, delta_epsilon [cap D(alpha).

In the sequel beta and gamma are two maximal monotone graphs on R verifying 0 in beta(0), 0 in gamma(0). It is well known that for every u in H there exists a unique function Pu in W^{2,2}(Omega) satisfying

(3) { -Delta Pu + gamma(Pu) + Pu delta u, a.e. in Omega; - (partial Pu / partial n) in beta(Pu), a.e. on Gamma,

where partial / partial n is the outward normal derivative; moreover || Pu ||_{2,2} <= c || u ||. We denote by c constants depending only on Omega, n, beta and gamma.

In [1] we have introduced a method that applies, in particular, to the study of the differentiability of the Green's operator P^(1). More precisely

(*) Instituto de Física e Matemática (Lisbon).

(**) Nella seduta del 12 aprile 1975.

(1) In [1] we derive from this result a theorem on the bifurcation points for the operator P.

we have proved that:

THEOREM I: (i) *If γ is differentiable at the origin with $\gamma' = +\infty$ then the operator P is Fréchet differentiable and $DP(0) = 0$.*

(ii) *If β and γ are differentiable at the origin with $\gamma' < +\infty$ and $\beta' = +\infty$ then the operator P is Fréchet differentiable at the origin and $DP(0) = A$ is the Green's operator for the linear Dirichlet problem*

$$(4) \quad \begin{cases} -\Delta Au + Au + \gamma' Au = u & \text{in } \Omega, \\ Au = 0 & \text{on } \Gamma. \end{cases}$$

(iii) *If $\gamma' < +\infty$ and $\beta' < +\infty$ then $DP(0) = A$ is the Green's operator for the linear problem*

$$(5) \quad \begin{cases} -\Delta Au + Au + \gamma' Au = u & \text{in } \Omega, \\ -\partial Au / \partial n = \beta' Au & \text{on } \Gamma. \end{cases}$$

Obviously Theorem I is equivalent to prove that

$$(6) \quad \lim_{\|u\| \rightarrow 0} \frac{\|Pu - Au\|}{\|u\|} = 0,$$

where $A = 0$ in case (i).

It was remarked to the author (oral communication) by J. Hernandez that one can prove that

$$(7) \quad \lim_{\|u\| \rightarrow 0} \frac{\|Pu - Au\|_{1,2}}{\|u\|} = 0.$$

The aim of this note is to verify that (7) is a trivial consequence of the estimates obtained in [1]. For brevity when we write " $\leq c\varepsilon \|u\|$ " it is understood that the corresponding estimate is true for $\|u\|$ sufficiently small.

Cases (ii) and (iii):

Put $Ru = Pu - Au$. In [1] we succeed in proving that (cf. [1], (1.21), (1.22) and (1.25))

$$(8) \quad \|\Delta Ru - Ru - \gamma' Ru\|_q \leq c\varepsilon \|u\| \quad \text{in cases (ii) and (iii),}$$

with $q < 2^{(2)}$, and we also show

$$(9) \quad \|Ru\|_2 \leq c\varepsilon \|u\| \quad \text{in case (ii),}$$

$$(10) \quad \|\partial Ru / \partial n + \beta' Ru\|_2 \leq c\varepsilon \|u\| \quad \text{in case (iii).}$$

(2) To prove (6) we had chosen in [1] $q = (2^*)'$ where $2^* = 2n/(n-2)$ is the Sobolev imbedding exponent of $W^{1,2}(\Omega)$ and $1/(2^*)' = 1 - (1/2^*)$. To prove (7) we made the same choice of q . If $n \leq 2$ then $q < 2$ can be arbitrarily; but in this case the results can be strengthened.

From (8), (9), (10) and from well known estimates for solutions of linear equations we deduce immediatly, in our paper [1], relation (6). But from exactly the same estimates (8), (9), (10) one trivially derives relation (7). In fact multiplying $-\Delta Ru + Ru + \gamma' Ru$ by Ru , integrating in Ω and applying Green's formulae it follows that (we recall that $\gamma' \geq 0$)

$$(11) \quad \|Ru\|_{1,2}^2 \leq \int_{\Gamma} (\partial Ru / \partial n) Ru \, d\Gamma + \int_{\Omega} (-\Delta Ru + Ru + \gamma' Ru) Ru \, dx.$$

In case (ii) from the corresponding estimates (8), (9) it follows then that

$$(12) \quad \|Ru\|_{1,2} \leq c\varepsilon \|u\|,$$

because $\| \cdot \|_{2^*} \leq c \| \cdot \|_{1,2}$.

Analogously in case (iii) the corresponding estimates (8), (10) give (12) because we have in (11)

$$\int_{\Gamma} (\partial Ru / \partial n) Ru \, d\Gamma \leq \int_{\Gamma} (\partial Ru / \partial n + \beta' Ru) Ru \, d\Gamma.$$

Finally (i) follows from (6) (i.e. from $\|Pu\| \leq c\varepsilon \|u\|$) and from equation (3).

We remark that (7) can be further generalized. Consider for example case (iii). Formula (8) holds for all $q < 2$ (as proved in [1]) and consequently (8) holds with $\| \cdot \|_q$ replaced by $\| \cdot \|_{s,2}$, for all $s < 0$. From this estimate, from (10) and from known results for linear equations it follows that $\|Ru\|_{3/2,2} \leq c\varepsilon \|u\|$; consequently

$$(13) \quad \lim_{\|u\| \rightarrow 0} \frac{\|Pu - Au\|_{3/2,2}}{\|u\|} = 0.$$

This relation can be generalized.

REFERENCES

- [1] H. BEIRÃO DA VEIGA - *Differentiability for Green's operators of variational inequalities and applications to the calculus of bifurcation points* (to appear in the «Journal of Functional Analysis»).

